This article was downloaded by: [New York University]

On: 22 May 2015, At: 04:22 Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House,

37-41 Mortimer Street, London W1T 3JH, UK



Communicative & Integrative Biology

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/kcib20

Pollinator shortage and global crop yield

Lucas A. Garibaldi^a, Marcelo A. Aizen^b, Saul Cunningham^c & Alexandra M. Klein^d

- ^a Universidad Nacional del Comahue
- ^b Universidad de Buenos Aires
- ^c CSIRO Entomology
- ^d University of California, Berkley; and University of Goettingen Published online: 01 Feb 2009.

To cite this article: Lucas A. Garibaldi, Marcelo A. Aizen, Saul Cunningham & Alexandra M. Klein (2009) Pollinator shortage and global crop yield, Communicative & Integrative Biology, 2:1, 37-39, DOI: 10.4161/cib.2.1.7425

To link to this article: http://dx.doi.org/10.4161/cib.2.1.7425

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

Article Addendum

Pollinator shortage and global crop yield

Looking at the whole spectrum of pollinator dependency

Lucas A. Garibaldi, 1,2,* Marcelo A. Aizen, 1 Saul A. Cunningham3 and Alexandra M. Klein4,5

¹Laboratorio Ecotono; INIBIOMA-CONICET and Centro Regional Bariloche; Universidad Nacional del Comahue; Río Negro, Argentina; ²Cátedra de Métodos Cuantitativos Aplicados; Facultad de Agronomía; Universidad de Buenos Aires; Buenos Aires, Argentina; ³CSIRO Entomology; Canberra Australia; ⁴Environmental Sciences Policy and Management; University of California, Berkeley; Berkeley, California USA; ⁵Agroecology; University of Goettingen, Germany

Abbreviations: FAO, food and agricultural organization of the united nations

Key words: agricultural production, developed world, developing world, ecosystem services, environmental degradation, FAO, pollination crisis

A pollinator decline caused by environmental degradation might be compromising the production of pollinator-dependent crops. In a recent article, we compared 45 year series (1961–2006) in yield, production and cultivated area of pollinator-dependent and nondependent crop around the world. If pollinator shortage is occurring globally, we expected a lower annual growth rate in yield for pollinator-dependent than nondependent crops, but a higher growth in cultivated area to compensate the lower yield. We have found little evidence for the first "yield" prediction but strong evidence for the second "area" prediction. Here, we present an additional analysis to show that the first and second predictions are both supported for crops that vary in dependency levels from nondependent to moderate dependence (i.e., up to 65% average yield reduction without pollinators). However, those crops for which animal pollination is essential (i.e., 95% average yield reduction without pollinators) showed higher growth in yield and lower expansion in area than expected in a pollination shortage scenario. We propose that pollination management for highly pollinator-dependent crops, such us renting hives or hand pollination, might have compensated for pollinator limitation of yield.

Meeting the growing demand in the amount and diversity of food, while dealing with increasing environmental degradation, is a major challenge for humanity. As a result of anthropogenic activities, managed and wild pollinators are declining in many regions of the world.¹⁻⁷ This could have important consequences for our food

*Correspondence to: Lucas A. Garibaldi; Laboratorio Ecotono; INIBIOMA-CONICET and Centro Regional Bariloche; Universidad Nacional del Comahue; Quintral 1250; 8400 Bariloche; Río Negro, Argentina; Email: garibald@agro.uba.ar

Submitted: 11/13/08; Accepted: 11/14/08

Previously published online as a *Communicative & Integrative Biology* E-publication: http://www.landesbioscience.com/journals/cib/article/7425

Addendum to: Aizen MA, Garibaldi LA, Cunningham SA, Klein AM. Long-term global trends in crop yield and production reveal no current pollination shortage but increasing pollinator dependency. Curr Biol 2008; 18:1572–5; PMID: 18926704; DOI: 10.1016/j.cub.2008.08.066.

supply, because approximately three-quarters of the crops cultivated worldwide depend to some degree on pollinators to produce seeds, fruits and vegetables.⁸⁻¹⁰ Indeed, pollinators promote either yield quantity or quality for crops; even those with a long history of artificial selection and/or more recently genetically engineering, such as canola, soybean and sunflower.¹¹⁻¹³ There is evidence at the local scale that diminished pollinator populations due to environmental degradation are limiting crop yield, ^{12,14-17} but only recently have we assessed if this phenomenon affects global food production.

In a recent paper, ¹⁸ we tested the hypothesis that a reduction in pollinator abundance is limiting crop yield at the global scale. We compared 45 year series (1961–2006) in yield, production and cultivated area of pollinator-dependent and nondependent crops. ⁸ These temporal trends could differ between the developed and developing world because of differences in agricultural intensification, and socioeconomic and environmental conditions. ¹⁹⁻²² We found that crop yield (Mt/ha) has increased on average by 1.5% per year from the values shown in 1961. This trend was similar for pollinator-dependent and nondependent crops in either the developed or developing worlds, thus providing no evidence that pollinator decline has yet affected crop yield at a global scale.

However, there are reasons to expect that the pollinator shortage may have a global impact in the near future. First, we found that the area devoted to pollinator-dependent crops has increased faster than that of nondependent crops in both the developed and developing worlds. ¹⁸ Therefore, the demand for animal pollination service is rising at the same time that pollinator abundance and diversity are declining. Indeed, if a pollinator shortage is limiting crop yield, a higher growth in area in pollinator-dependent crops is expected to maintain production rates. Second, we have found a trend of lower growth rate in yield in crops that depend highly on pollinators compared to those with low or none dependence. ¹⁸ All these results could be interpreted as an early warning sign of global pollinator declines effects.

Estimation of pollinator dependency effects on global crop production must recognize that the degree of crop dependence varies greatly, such that absence of pollinators would reduce yield by 100% in some crops, and by only a few percent for other crops. ^{9,11} In our recent paper, we classified crops into three categories of pollinator

dependency: no dependence, low dependence and high dependence. ¹⁸ Here, our analysis classify crops into five categories of pollinator dependency based on Klein et al. ⁸ We also provide new estimates of pollinator dependency effects by regression analyses of the relationship between crops annual growth in relative yield, and cultivated area, with their level of pollinator dependency.

We used the dataset described by Aizen et al., 18 and compiled annual data on yield and cultivated area for 87 crops during 1961-2006 for the developed and developing worlds from the FAO statistics.²³ These crops collectively accounted for 83% of total global food production during 2006. Using information from pollinator-exclusion experiments provided by Klein et al. review,8 we classified each crop into one of five categories according to the average reduction in yield due to the absence of pollinators: 0% (no dependence), 5%, 25%, 65% and 95%. The nondependent category included crops pollinated abiotically (wind) or autogamously, and crops that depend on pollinators but are cultivated for the vegetative parts (e.g., leaves, stems, tubers). For each crop in each region, we first standardized the change in each variable x (i.e., yield or area) during year t relative to its value during 1961 as Δx_t = $100 \cdot (x_t - x_{1961})/x_{1961}$. We then estimated the slope,

 β , of the linear relation between Δx_t and year (i.e., %/yr) as an unbiased estimate of the linear average annual growth in the relative yield or area, despite temporal autocorrelation. For example, a slope of 2 for area indicates that the cultivated area of a given crop increased, on average, by 2%/yr in relation to the area cultivated in 1961. For these analyses we excluded crops that were not cultivated in 1961 as explained in Aizen et al. 18

Pollinator shortfalls would produce lower annual growth in yield in crops with greater dependency on pollinators. We also expect a higher annual growth in area in crops with greater dependency on pollinators to compensate the lower yields. In both the developed and developing worlds, we found evidence that support these predictions for the crops that range in dependency levels from 0 to 65% (Fig. 1). However, those crops for which animal pollination is essential (i.e., 95% yield reduction without pollinators) showed higher growth in yield and lower growth in area than expected by a pollination shortage scenario (Fig. 1).

Pollination management in these highly dependent crops might be decreasing the pollinator limitation on yield produced by environmental degradation. Brazil nuts showed a negative growth in yield but cultivated area did not rise as fast as expected (Table 1). This is reasonable because Brazil nut trees are not commercially planted; the nuts are collected within the Amazonian rainforest that has been decreasing in area because of deforestation. Watermelon, melon and pumpkin (all vines in the family Cucurbitaceae) are frequently pollinated by managed honeybees in large-scale^{25,26} or by hand in small-scale production systems (authors' personal observation). Cocoa trees are managed in agroforestry systems which usually provide resources for their pollinating midges.^{27,28} Finally, vanilla is commercially grown only under hand pollination management techniques.²⁹ Therefore, we propose that there is an interaction between




Figure 1. Mean ± 1 SE of the annual relative growth in yield and cultivated area for crops differing in their pollinator dependence. The lines are linear regressions without considering the 95% dependence group and taking each crop as a replicate. In parenthesis is the number of crops within each group.

the pollinator-dependence spectrum and management responses. At the low end of the range, where pollinators make less difference in yield, management will commonly ignore pollinators. At the high end of the range, where crops will fail or be hugely diminished in the absence of pollinators, most growers will actively manage their pollination.

Acknowledgements

We thank Sofía Gonzalez for useful comments and suggestions. This work was conducted partly within the framework provided by the Restoring Pollination Services Working Group supported by the National Center for Ecological Analysis and Synthesis, a Center funded by NSF (grant no. DEB-0072909). We acknowledged additional funding by the Argentina National Council for Research (PIP 5066) and the National University of Comahue (B126/04). Lucas A. Garibaldi holds a fellowship from the Argentina National Council for Research and Marcelo A. Aizen is a career researcher of the same agency. Alexandra M. Klein is supported by the Alexander von Humboldt Foundation.

References

- Biesmeijer JC, Roberts SPM, Reemer M, Ohlemuller R, Edwards M, Peeters T, et al. Parallel declines in pollinators and insect pollinated plants in Britain and the Netherlands. Science 2006: 313:351-4.
- Kluser S, Peduzzi P. Global Pollinator Decline: A Literature Review. Geneva: UNEP/GRID 2007.
- National Research Council. Status of Pollinators in North America. Washington DC: National Academies Press 2007.
- 4. Oldroyd BP. What's killing American honey bees? PLoS Biol 2007; 5:168.
- Fitzpatrick U, Murray TE, Paxton RJ, Breen J, Cotton D, Santorum V, et al. Rarity and decline in bumblebees—a test of causes and correlates in the Irish fauna. Biol Conserv 2007; 136:185-94.
- Kosior A, Celary W, Olejniczak P, Fijal J, Kro'l W, Solarz W, et al. The decline of the bumble bees and cuckoo bees (Hymenoptera: Apidae: Bombini) of western and central Europe. Orvx 2007; 41:79-88.
- 7. Watanabe ME. Pollination worries rise as honey bees decline. Science 1994; 265:1170

Table 1 Annual growth in yield ($\beta_{\Delta yield}$) and cultivated area ($\beta_{\Delta area}$) during 1961–2006 for highly pollinator-dependent crops (i.e., 95% average reduction in yield in the absence of pollinators) in the developed and developing world

Species	Сгор	Developed world ¹		Developing world	
	•	$\beta_{\Delta yield}$ (%/yr)	$eta_{\Delta lpha rea}$ (%/yr)	$eta_{\Delta extsf{yield}}$ (%/yr)	$\beta_{\Delta area}$ (%/yr)
Bertholletia excelsa	Brazil nut, Para nut, Cream nut	- -	-	-0.144	1.072
Citrullus lanatus	Watermelon	3.003	-0.921	3.814	3.635
Cucumis melo	Cantaloupe, Melon	1.935	0.554	1.895	3.369
Cucurbita maxima, C. mixta, C. moschata, C. pepo	Pumpkin, Squash, Gourd, Marrow, Zucchini	16.775	-1.150	0.945	5.103
Theobroma cacao	Cocoa	-	-	2.085	1.690
Vanilla planifolia, V. pompona	Vanilla	-	-	1.149	6.135

¹Brazil nut, cocoa and vanilla are not commonly cultivated in the developed world and thus do not appear in the FAO statistics

- Klein AM, Vaissie're BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, et al. Importance of pollinators in changing landscapes for world crops. Proc R Soc B 2007; 274:303-13.
- Roubik DW. Pollination of cultivated plants in the tropics. FAO Agricultural Services Bulletin 118. Rome:FAO 1995.
- Williams IH. The dependence of crop production within the European Union on pollination by honey bees. Agricultural Zoology Reviews 1994; 6:229-57.
- Chiari WC, Toledo VAA, Ruvolo-Takasusuki MCC, Oliveira AJB, Sakaguti ES, Attencia VM, et al. Pollination of soybean (Glycine max L. Merril) by honeybees (*Apis mellifera* L.). Brazilian Archives of Biology and Technology 2005; 48:31-6.
- Greenleaf SS, Kremen C. Wild bees enhance honey bees' pollination of hybrid sunflower. Proc Natl Acad Sci USA 2006; 103:13890-5.
- Morandin LA, Winston ML. Pollinators provide economic incentive to preserve natural land in agroecosystems. Agric Ecosyst Environ 2006; 116:289-92.
- Priess JA, Mimler M, Klein A, Schwarze S, Tscharntke T, Steffan-Dewenter I. Linking deforestation scenarios to pollination services and economic returns in coffee agroforestry systems. Ecological Applications 2007; 17:407-17.
- Blanche KR, Ludwig JA, Cunningham SA. Proximity to rainforest enhances pollination and fruit set in orchards. J Appl Ecol 2006; 43:1182-7.
- Ricketts TH, Daily GC, Ehrlich PR, Michener CD. Economic value of tropical forest to coffee production. Proc Natl Acad Sci USA 2004; 101:12579-82.
- Richards AJ. Does low biodiversity resulting from modern agricultural practice affect crop pollination and yield? Ann Bot (Lond) 2001; 88:165-72.
- Aizen MA, Garibaldi LA, Cunningham SA, Klein AM. Long-term global trends in crop yield and production reveal no current pollination shortage but increasing pollinator dependency. Curr Biol 2008; 18:1-4.
- Matson PA, Parton WJ, Power AG, Swift MJ. Agricultural intensification and ecosystem properties. Science 1997; 277:504-9.
- Allen JC, Barnes DF. The causes of deforestation in developing countries. Annals of the Association of American Geographers 1985; 75:163-84.
- Conway G. The doubly green revolution: Balancing food, poverty and environmental needs in the 21st century. In: Lee DR, Barrett CB, eds. Tradeoffs or Synergies? Agricultural Intensification, Economic Development and the Environment. Wallinford: CABI Publishing 2001; 17-34.
- Evenson RE, Gollin D. Assessing the impact of the green revolution, 1960 to 2000. Science 2003; 300:758-62.
- FAOSTAT. Data available at http://faostat.fao.org/default.aspx Last accessed in January 2008.
- 24. Murtaugh PA. Simplicity and complexity in ecological data analysis. Ecology 2007; 88:56-62.
- Morse RA, Calderone NW. The value of honey bees as pollinators of U.S. crops in 2000. Bee Culture 2000; 2-15.
- Walters SA. Honey bee pollination requirements for triploid watermelon. HortScience 2005; 40:1268-70.
- Bos MM, Steffan-Dewenter I, Tscharntke T. Shade tree management affects fruit abortion, insect pests and pathogens of cacao. Agric Ecosyst Environ 2007; 120:201-5.
- Young AM. Effects of shade cover and availability of midge breeding sites on pollinating midge populations and fruit set in two cacoa farms. J Appl Ecol 1982; 19:47-63.
- Davis EW. Experiences with growing vanilla (Vanilla planifolia). Acta Horticulturae 1983; 132:23-9.

39