




ELSEVIER

Contents lists available at ScienceDirect

Cretaceous Research

journal homepage: www.elsevier.com/locate/CretRes

Upper Maastrichtian and Danian bryozoans from Northern Patagonia, Argentina

Soledad S. Brezina^{a, *}, Paul D. Taylor^b, María V. Romero^{c, d}, Evangelina E. Palópolo^{a, c}, Silvio Casadío^{a, c}

^a Universidad Nacional de Río Negro, Instituto de Investigación en Paleobiología y Geología, Av. Roca 1242, General Roca, Río Negro, R8332FDJ, Argentina

^b Department of Earth Sciences, Natural History Museum, Cromwell Road, London, SW7 5BD, UK

^c Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina

^d Estación Costera J.J. Nágera, Instituto de Investigaciones Marinas y Costeras, Rodríguez Peña 4046, B7602GSD, Mar del Plata, Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 12 October 2020

Received in revised form

4 April 2021

Accepted in revised form 4 April 2021

Available online 20 April 2021

Keywords:

Taxonomy

Cyclostomata

Cheilostomata

Maastrichtian

Danian

Neuquén Basin

Patagonia

ABSTRACT

The specimen-rich and diverse bryozoan fauna encrusting oyster shells from the Maastrichtian and Danian Jagüel and Roca formations in the Neuquén Basin (Patagonia, Argentina) is described. Thirteen cyclostome species are recorded, along with 18 cheilostomes. We introduce two new species: *Akatopora kaufmanni* sp. nov. and *Eoporella lunata* gen. et sp. nov. The latter is a homeomorph of the common Miocene–Recent genus *Microporella*. Ten of the cyclostomes and ten of the cheilostomes could not be identified at species-level because of preservational limitations or lack of diagnostic characters in the available specimens. The most common bryozoans present are sheet-like colonies, among which the ascophoran cheilostome *Balantiostoma* is particularly well represented. Results show that in northern Patagonia the diversity of encrusting bryozoans associated with oyster shells exhibits no major changes across the K/Pg boundary. However, an important increase in the diversity is recorded during the upper Danian.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

The fossil record of bryozoans is rich and extends back to the Ordovician. Bryozoans radiated rapidly during the Late Cretaceous and were diverse and abundant in many marine environments, peaking for both cheilostomes and cyclostomes in the Maastrichtian prior to the K/Pg boundary (Taylor and Waeschenbach, 2015). Late Cretaceous bryozoans are best known from northern Europe thanks largely to the works of the prolific taxonomist Erhard Voigt (1905–2004). Rich bryozoan faunas also occur in the Late Cretaceous of central Asia and Europe (Voigt, 1967; Favorskaya, 1985, 1987, 1988, 1992; Koromyslova et al., 2018a, b, c; Koromyslova et al., 2019a, b; Koromyslova and Seltser, 2020), southeastern North America (Taylor and McKinney, 2006; Sogot et al., 2013; 2014), Madagascar (Di Martino et al., 2018) and India (Guha and Nathan, 1996; Taylor and Di Martino, 2018). Bryozoan assemblages reported from tropical environments of Late Cretaceous age are

scarce (Di Martino and Taylor, 2013). Late Cretaceous and Paleocene bryozoans are poorly known from South America, where knowledge is largely limited to the work by Canu (1911). Many of the species described by Canu have never been revised and are difficult to interpret from his figures which are retouched photographs. Furthermore, several specimens in his collection housed at the Museo Argentino de Ciencias Naturales Bernardino Rivadavia are missing. This hampers identification of species, which is a strong impediment for comparisons with coeval associations and for studies that could reveal how bryozoans responded to changes associated with the Cretaceous–Paleogene boundary events in South America.

Bryozoans commonly occur as encrustations on hard substrates that include rocks, clasts and the skeletons of living and dead organisms (Taylor and Wilson, 2003). Prominent among skeletal substrates are oysters, which have a high preservation potential in the fossil record. Oysters from the Late Cretaceous–early Paleogene marine successions in Patagonia are abundant and quite well-preserved (Casadío, 1998). They support a rich variety of boring and encrusting sclerobionts, including sponges, polychaetes, bivalves, fungi, algae, barnacles and bryozoans (Brezina et al., 2014,

* Corresponding author.

E-mail address: sbrezina@unrn.edu.ar (S.S. Brezina).

2017). The aim of our work here is to describe the Maastrichtian and Danian bryozoan faunas encrusting oyster shells from the Jagüel and Roca formations of the Neuquén Basin (Patagonia, Argentina). Aside from its regional significance, this study adds to our limited knowledge of Southern Hemisphere fossil bryozoan faunas from the Late Cretaceous (Taylor, 2019), and to extinction and survival of bryozoans across the K/Pg boundary (e.g. Stilwell and Håkansson, 2012).

2. Geological setting

The Neuquén Basin in west-central Argentina covers parts of the provinces of Río Negro, Neuquén, La Pampa and Mendoza (Fig. 1). Rocks exposed in the basin range in age from Late Triassic to Paleogene. A major transgression from the South Atlantic into the basin occurred during the Maastrichtian to Danian, a time of relative tectonic quiescence and low volcanic activity (Malumián and Nández, 2011). This transgression reduced the southern tip of South America (Patagonia) to an archipelago during the K/Pg boundary (Aguirre-Urreta et al., 2008). In the Neuquén Basin, the sediments deposited by this transgression are represented by the Jagüel and Roca formations.

In the deeper parts of the basin mudstones dominate the Maastrichtian and lower Danian deposits (Jagüel Formation). They are massive and presumably intensely bioturbated, indicating a well-oxygenated seafloor (Scasso et al., 2005). The lithology and fossils indicate deposition at mid-shelf depths (Brezina et al., 2014). Shallower proximal areas mainly consist of limestones of the Roca

Formation. This formation, which comprises bioclastic packstones and grainstones deposited in subtidal to intertidal settings, shows a regressive character, and has been dated as Maastrichtian in the northern part of the basin (north of 36°S) but is Danian in age for outcrops in the central and southern parts.

Circulation within the Neuquén Basin was slightly restricted during the upper Maastrichtian and lower Danian (Scasso et al., 2005). Evaporites and mixed carbonate-siliciclastic lithologies in the northernmost part of the Neuquén Basin suggest semi-arid and hypersaline conditions in that part of the basin (Kiesling et al., 2006). The occurrence of salinity stratification in the north-western part of the Neuquén Basin during the upper Maastrichtian is indicated by high percentages of prasinophytes that suggest stratified and saline waters (Prámparo et al., 2014). Foraminiferal and nannofossil analyses in the central part of the basin indicate that sedimentation during the upper Maastrichtian took place in relatively shallow middle neritic depths (Keller et al., 2007) and normal marine conditions, evidenced by the presence of relatively few terrestrial palynomorphs (Woelders et al., 2017; Guler et al., 2019).

The TEX₈₆ based sea surface temperatures obtained by Woelders et al. (2017) in the Jagüel and Roca formations show that the last two million years of the Cretaceous were characterized by multiple warming and cooling phases, with average sea surface temperatures varying between 23 °C and 29 °C. Within the K/Pg boundary layer, sea surface temperature dropped to <19 °C, followed by a rapid warming to 31.6 °C. de Winter et al. (2018), based on stable oxygen isotope thermometry of Maastrichtian oyster shells,




Fig. 1. Map of fossiliferous localities from which the material comes from. BdJ: Bajada del Jagüel; CBA: Cerros Bayos; CBU: Cerro Butaló; CdP: Casa de Piedra; GR: General Roca; Hu: Huantraico; LM: Liu Malal; RC: Ranquil-Có.

retrieved water temperatures of 11 °C. Discrepancy with TEX₈₆ palaeothermometry was attributed to seasonal bias in the growth of the oysters, while TEX₈₆ data appear to be biased towards warmer sea surface water temperatures (de Winter et al., 2018).

3. Material and methods

All of the bryozoans described here are encrusting species. The oyster shells to which they are cemented were collected in the field by bulk sampling from eight localities (Fig. 2). Cleaning methods included scrubbing under running water with a soft toothbrush and brief ultrasonic cleaning. The bryozoans were sorted and preliminary identifications made with the aid of binocular microscopes. The best-preserved specimens were selected for uncoated scanning electron microscopy (SEM) using a LEO 1455VP operating in low vacuum mode at the Natural History Museum, London. Back-scattered images were captured digitally.

Existing material in museum and other collections, as well as specimens newly collected in the field or described in a recent paper (Taylor and Brezina, 2018), were utilized for this study. Repository abbreviations for material in museum collections are as

follows: GHUNLPam, Cátedra de Geología Histórica de la Universidad Nacional de La Pampa, Santa Rosa, Argentina; MACN, Museo Argentino de Ciencias Naturales, Ciudad Autónoma de Buenos Aires, Argentina; MPEF, Museo Paleontológico Egidio Feruglio, Trelew, Argentina; NHMUK, Natural History Museum, London, UK.

Except where noted, measurements were made from zones of astogenetic repetition. All zooidal measurements were taken from digital SEM images. Each measurement is given in the text as mean ± standard deviation, with the number of specimens used and total number of measurements or counts made enclosed in parenthesis.

Measurements of the cheilostomes use the following abbreviations: AL, avicularium length; AW, avicularium width; OL, orifice length; OW, orifice width; OOW, ovicellate zooid orifice width (where different from normal orifice width); OpL, opesia length; OpW, opesia width; OvL, ovicell length; OvW, ovicell width; ZL, autozooid length (as seen on colony surface); ZW, autozooid width (as seen on colony surface).

Measurements of cyclostomes are identified by some of the same abbreviations as for cheilostomes but with the addition of




Fig. 2. Selected stratigraphic sections, illustrating lithology and marking beds that yielded oysters with bryozoans. Dashed line indicates K-Pg boundary.

the following: AD, diameter of equidimensional apertures; AS, distance between midpoints of adjacent apertures; ASW distance between midpoints of adjacent apertures within a row or fascicle; BCL, brood chamber length; BCW, brood chamber width; BrD, ADmn, minimum diameter of apertures; ADmx, maximum diameter of apertures; FS, Distance between centres of successive fascicles; branch diameter or width; FWL, frontal wall length of single zooid; FWW, maximum frontal wall width of single zooid; Gap, distance between edges of adjacent fascicles; GL, gonozooid length including proximal portion and brood chamber; GW, gonozooid width; NAD, nanozooid aperture diameter; OD, oocypore diameter.

4. Systematic palaeontology

All the bryozoans described in this work encrust oyster shells from the upper Maastrichtian to the upper Danian of Northern Patagonia, and are listed in Table 1.

Order Cyclostomata Busk, 1852

Suborder Tubuliporina Milne Edwards, 1838

Family Stomatoporidae Pergens and Meunier, 1886

Genus *Voigttopora* Bassler, 1952

Type species: *Alecto calypso* d'Orbigny, 1849, 'Senonian' [probably Santonian], Saintes, Charente Maritime, France (see Illies, 1976).

Voigttopora sp.

Fig. 3A–C

Material. MPEF-PI 6132.4, encrusting colony on a valve of *Cubitostrea ameghinoi* Ihering, 1902; upper Danian, Roca Formation, Cerros Bayos, La Pampa.

Description. Colony encrusting, uniserial (Fig. 3A, B), branches originating solely by lateral budding at 70°–90° to the parent branch (Fig. 3C); up to at least 6 autozooids between successive lateral branches. Autozooids broad, parallel-sided, their frontal walls containing abundant circular pseudopores; preserved peristomes short, apertures circular. Ancestrula not observed.

Measurements. AS 231 ± 69.15 µm (2, 13); AD 107 ± 13.34 µm (1, 3).

Remarks. This species somewhat resembles *Voigttopora maconensis* Taylor and McKinney, 2006 from the Late Cretaceous of the southeastern USA, but has shorter, stouter autozooids. Another North American species, *V. thurni* Taylor and McKinney, 2006, possesses autozooids of similar length to those of the Roca *Voigttopora* but ramifications occur both dichotomously and by lateral branching (as in *V. calypso*, the type species of *Voigttopora*), whereas the Roca species shows only lateral ramifications. The lack of an ancestrula and early astogenetic stages in available material discourages the introduction of a new species for the Roca material.

Family Oncousoeciidae Canu, 1918

Genus *Oncousoecia* Canu, 1918

Table 1

List of encrusting bryozoan species associated with oysters recorded across the K/Pg boundary.

Bryozoan species	Colony form	Collection number
Upper Maastrichtian		
Cheilostomata		
<i>Flustrellaria</i> sp.	encrusting, multiserial	MPEF-PI 6132.21
? <i>Akatopora</i> sp. 1	encrusting, multiserial	MPEF-PI 6132.20
<i>Poricella tripora</i>	encrusting, multiserial	MPEF-PI 6132.67
? <i>Trichinopolia</i> sp.	encrusting, multiserial	MPEF-PI 6132.20
Lower Danian		
Cheilostomata		
<i>Conopeum okaiana</i>	encrusting, multiserial	GHUNLPam 17423
<i>Aspidostoma onychocecliferum</i>	encrusting, multiserial	MPEF-PI 6132.68
<i>Tricephalopora</i> sp. 1	encrusting, multiserial	MPEF-PI 6132.19
Upper Danian		
Cyclostomata		
<i>Voigttopora</i> sp.	encrusting, uniserial	MPEF-PI 6132.4
<i>Oncousoecia?</i> sp.1	encrusting, oligoserial	MPEF-PI 6132.25
<i>Oncousoecia?</i> sp. 2	encrusting, oligoserial	MPEF-PI 6132.9
<i>Oncousoecia?</i> cf. <i>striata</i>	encrusting, oligoserial	MPEF-PI 6132.26
<i>Axilosoezia giselae</i>	encrusting, uniserial	MPEF-PI 6132.28
<i>Platonea</i> sp.	encrusting, oligoserial	MPEF-PI 6132.5
<i>Plagioecia</i> aff. <i>Cristata</i>	encrusting, multiserial	MPEF-PI 6132.18
<i>Mesenteripora</i> sp.	encrusting, multiserial	MPEF-PI 6132.28
<i>Actinopora robertsoniana</i>	encrusting, multiserial	MPEF-PI 6132.17
' <i>Berenicea</i> ' sp.	encrusting, multiserial	MPEF-PI 6132.13
<i>Disporella?</i> <i>discoidea</i>	encrusting, multiserial	MPEF-PI 6132.65
<i>Ceripora</i> sp.	encrusting, multiserial	MPEF-PI 6132.2
Cheilostomata		
<i>Electra</i> sp.	encrusting, multiserial	MPEF-PI 6132.1
? <i>Labioporella</i> sp.	encrusting, multiserial	MPEF-PI 6132.8
<i>Pyriporella ameghinoi</i>	encrusting, multiserial	MPEF-PI 6132.18
<i>Akatopora kaufmanni</i> sp. nov.	encrusting, multiserial	NHMUK D32352a
<i>Cianotremella gigantea</i>	encrusting, multiserial	NHMUK D32352b
<i>Monoporella convexa</i>	encrusting, multiserial	MPEF-PI 6132.15–16
<i>Tricephalopora?</i> sp.	encrusting, multiserial	MPEF-PI 6132.3
<i>Poricella</i> sp.	encrusting, multiserial	MPEF-PI 6132.23
<i>Tremogasterina problematica</i>	encrusting, multiserial	MPEF-PI 6132.11
<i>Balantiostoma spectabilis</i>	encrusting, multiserial	MPEF-PI 6132.15–66
<i>Balantiostoma elongate</i>	encrusting, multiserial	MPEF-PI 6132.6–12
<i>Balantiostoma</i> sp.	encrusting, multiserial	MPEF-PI 6132.7–13–14
<i>Eoporella lunata</i> gen. et sp. nov.	encrusting, multiserial	MPEF-PI 6132.6

Type species: *Tubulipora lobulata* Canu, 1918, Recent, British Isles (see Taylor and Zatoń, 2008).

?*Oncousoecia* sp. 1

Fig. 3D–E

Material. MPEF-PI 6132.25, encrusting colony on a valve of *Pycnodonte* (*Phygraea*) *sarmientoi* Casadío, 1998; upper Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colony encrusting, oligoserial, branches smoothly lobate. Autozooids elongate, the flat frontal walls perforated by subcircular to drop-shaped pseudopores 10 µm in diameter (Fig. 3D); zooidal boundaries not clearly defined; preserved peristomes short, tapering distally; apertures 50 µm in diameter (Fig. 3E). Kenozooids probably present along lateral edges of branches where autozooidal apertures are lacking. Gonozooids and early astogenetic stages not observed.

Measurements. AS 196 ± 48.70 µm (1, 18); AD 54 ± 12.54 µm (1, 19).

Remarks. Although this species is consistent with *Oncousoecia* in colony-form and autozooidal morphology, gonozooids have not been observed and its assignment to this genus must remain tentative. It also resembles, as does ?*Oncousoecia* sp. 2 described below, some species of the Cretaceous stomatopod *Proboscina* Pitt and Taylor, 1990, a genus lacking basal gonozooids.

?*Oncousoecia* sp. 2

Fig. 3F–H

Material. MPEF-PI 6132.9, colony encrusting a valve of *Pycnodonte* (*Phygraea*) *sarmientoi*; upper Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colony encrusting, oligoserial with bifurcating branches; branch surface marked by fine transverse rugae (Fig. 3F–G). Autozooids elongate with slightly convex frontal walls perforated by subcircular pseudopores 10 µm in diameter (Fig. 3H). Zooidal boundaries moderately well defined; preserved peristomes short, thick; apertures circular. Gonozooids and early astogenetic stages not observed.

Measurements. AS 387 ± 101.16 µm (1, 19); AD 83 ± 11.97 µm (1, 16); FWL 403 ± 43.17 µm (1, 8); FWW 160 ± 28.02 µm (1, 13).

Remarks. Despite the absence of gonozooids, the oligoserial colony-form allows tentative assignment of this species to *Oncousoecia*. The autozooids of ?*Oncousoecia* sp. 2 are appreciably larger than those of ?*O.* sp. 1 (c. 80 µm vs. c. 55 µm) and zooidal boundaries are more distinct because of the greater convexity of the frontal walls.

?*Oncousoecia* cf. *striata* Canu, 1911

Fig. 4A–F

cf. 1911 *Proboscina striata* Canu, p. 267, pl. 9, figs 8, 9.

Material. MPEF-PI 6132.26, colony encrusting a valve of *Pycnodonte* (*Phygraea*) *sarmientoi*; lower Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colonies encrusting, oligoserial with bifurcating branches 2–4 zooids wide (Fig. 4A). Ancestrula long, protoecium a flattened hemisphere with a concentric ridge inwards of the margin, without visible pseudopores. Two budded autozooids before the high-angled first branch bifurcation (Fig. 4B–C). Autozooids slightly convex with narrow salient boundary walls; apertures isolated or irregularly clustered into groupings of 2–5 (Fig. 4D, F). Preserved peristomes short to moderate in length. Possible base of an erect stem observed at distal end of an encrusting branch (Fig. 4E).

Measurements. AS 160 ± 76.78 µm (2, 16); AD 96 ± 11.11 µm (1, 9).

Remarks. Specimens studied here have somewhat higher branches than those of the material of ?*O. striata* (MACN-PI 1880) described by Canu (1911) and may represent a distinct species. The lack of gonozooids makes generic attribution difficult but colony-form

suggests that this species, as well as the species of Canu (1911), most likely belongs to *Oncousoecia*. However, it may alternatively be the encrusting base of a genus that normally grows erect, such as *Entalophoroecia* Harmelin, 1976.

Genus *Axilosoecia* Taylor and Brezina, 2018

Type species: *Axilosoecia giselae* Taylor and Brezina, 2018, upper Danian, Roca Formation, La Pampa, Argentina.

***Axilosoecia giselae* Taylor and Brezina, 2018**

Fig. 5A–E

2018 *Axilosoecia giselae* Taylor and Brezina, p. 442, fig. 1A–D.

Material. MPEF-PI 6132.28, 6132.51 (holotype), MPEF-PI 6132.28, 6132.53 (paratype) encrusting valves of the oyster *Cubitostrea ameghinoi*, upper Danian, Roca Formation, Cerros Bayos, La Pampa.

Description. Colony encrusting, uniserial, with branches bifurcating at about 90° (Fig. 5A), each internode consisting of one or occasionally two or three zooids (Fig. 5B). Early astogenetic stages with autozooids, about 200 µm long and higher angled bifurcations. Autozooids small, elongate, about 290–400 µm long by 90 µm wide in the zone of astogenetic repetition; apertures tiny, subcircular, about 0.04–0.05 mm wide, preserved peristomes short. Gonozooids positioned in axils of branch bifurcations, long proximal part indistinguishable from an autozooid, distal bulbous part seeming to originate from the peristome and descending onto the substrate (Fig. 5C–D), usually longitudinally elliptical, 230–400 µm long by 200–400 µm wide; oeciopore subterminal, small, transversely elliptical or subcircular, minute, about 20–30 µm wide (Fig. 5E).

Measurements. BrD 91 ± 15 µm (1, 5); GL 269 ± 27 µm (1, 2); GW 250 µm (1, 2).

Remarks. This was one of two new species of *Axilosoecia* described when the genus was introduced by Taylor and Brezina (2018). Both species are characterized by uniserial, runner-like colonies with gonozooids located in the axils of the branch bifurcations. The second species, *A. mediorubiensis* Taylor and Brezina, 2018 from the early Miocene of New Zealand, has longer autozooids, usually more zooids per internode, and gonozooids lacking an autozooid-like proximal part. Since the paper introducing *Axilosoecia* was published, an undescribed species referable to this genus has been found in collections from the Coon Creek Member (Ripley Formation) of New Albany, Union County, Mississippi (PDT unpublished). This Maastrichtian species more strongly resembles *A. giselae* than it does *A. mediorubiensis*.

Family Tubuliporidae Johnston, 1838

Genus *Platonea* Canu and Bassler, 1920

Type species: *Reptotubigera phillipsae* Harmer, 1915, Recent, Loyalty Island, Australia.

***Platonea* sp.**

Fig. 6A–B

Material. MPEF-PI 6132.5, colony encrusting a valve of *Pycnodonte* (*Phygraea*) *sarmientoi*; upper Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colony encrusting, comprising a single, curved lobate branch (Fig. 6A) with a wide crescentic growing edge. Ancestrula short, protoecium hemispherical, 95 µm in diameter, smooth surfaced, apparently lacking pseudopores, giving rise to a short distal tube (Fig. 6B). Autozooids in closely spaced, curved transverse rows on either side of branch midline, each row typically with 3 or 4 connate apertures rounded rectangular in shape. Frontal walls convex, the furrows between them delineating clearly the zooidal boundaries. A possible gonozooid with broken roof visible.

Measurements. ADmx 93 ± 5 (1, 8); ADmn 82 ± 4 (1, 6); FS 219 ± 13 µm (1, 4).




Fig. 3. **A–C**, *Voigtopora* sp., MPEF-PI 6132.4, upper Danian, Roca formation, Cerros Bayos, La Pampa. **A, B**, general view of two colonies (scale bars: 400 μ m). **C**, lateral branching (scale bar: 200 μ m). **D–E**, *?Oncousoecia* sp. 1., MPEF-PI 6132.25, upper Danian, Roca Formation, Casa de Piedra, La Pampa. **D**, Cylindrical peristomes with thin and elevated edges; the colony is overgrown by *Disporella* sp. to the left (scale bar: 200 μ m). **E**, autozooids walls perforated by subcircular to drop-shaped pseudopores (scale bar: 100 μ m). **F–H**, *?Oncousoecia* sp. 2, MPEF-PI 6132.9, upper Danian, Roca Formation, Casa de Piedra, La Pampa. **F**, general view of the colony (scale bar: 600 μ m). **G**, surface with fine concentric rugae parallel to the growing margin (scale bar: 400 μ m). **H**, frontal walls perforated by approximately subcircular pseudopores (scale bar: 200 μ m).

Remarks. This small colony may perhaps be the species incorrectly attributed by [Canu \(1911, p. 275\)](#) to *Idmonea carinata* [Römer, 1840](#). It is also similar to *Platonea adnata* [Taylor and McKinney, 2006](#) from the Maastrichtian Peedee Formation of North Carolina, which,

however, has larger autozooids and a protoecium of substantially greater diameter (165–196 μ m vs. 95 μ m in the Roca specimen). Introduction of a new species for the Roca cyclostome is deferred




Fig. 4. *?Oncousoecia* cf. *striata* ([Canu, 1911](#)), MPEF-PI 6132.26, lower Danian, Roca Formation, Casa de Piedra, La Pampa. **A**, general view of a colony (scale bar: 1 mm). **B**, ancestrula and three early generations of autozooids (scale bar: 300 μ m). **C**, detail of ancestrula and protoecium (scale bar: 100 μ m). **D**, autozooids clustered into groupings of two or three (scale bar: 200 μ m). **E**, possible broken erect branch (scale bar: 300 μ m). **F**, second colony with autozooidal apertures in groups of two or three (scale bar: 400 μ m).




Fig. 5. *Axilosocia giselae* Taylor and Brezina, 2018, MPEF-PI 6132.28, upper Danian, Roca Formation, Cerros Bayos, La Pampa. **A**, overview of a ramifying colony (scale bar: 1 mm). **B**, branch bifurcations, autozooids and two gonozooids (scale bar: 400 µm). **C**, two gonozooids showing bulbous brood chambers extending into the branch axils (scale bar: 200 µm). **D**, a pair of gonozooids (scale bar: 100 µm). **E**, gonozooid brood chamber and subterminal oocypore (scale bar: 100 µm).

pending the discovery of an intact gonozooid complete with oocypore.

Family Plagioeciidae Canu, 1918

Genus *Plagioecia* Canu, 1918

Type species: *Tubulipora patina* Lamarck, 1816, Recent, Europe.

Plagioecia sp.

Fig. 6C–F

Material. MPEF-PI 6132.18. Colony encrusting a valve of *Ostrea wilkensis*; upper Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, reniform in overall shape due to the non-preservation of one side (Fig. 6C), probably




Fig. 6. **A–B.** *Platonea* sp., MPEF-PI 6132.5, NP4 Biozone, upper Danian, Roca Formation, Casa de Piedra, La Pampa. **A**, general view of a colony with V-shaped fascicles, possible gonozooid indicated by an arrow (scale bar: 500 µm). **B**, detail of ancestrula and its hemispherical protoecium (scale bar: 100 µm). **C–F.** *Plagioecia* sp., MPEF-PI 6132.18, NP4 Biozone, upper Danian, Roca Formation, General Roca, Río Negro. **C**, general view of the colony (scale bar: 500 µm). **D–E**, two gonozooids from the same colony, with the oocypore indicated by an arrow in **E** (scale bars: 500 µm). **F**, detail of peristomes and pseudopores (scale bar: 100 µm).




Fig. 7. *Plagioecia* aff. *cristata* Taylor and McKinney, 2006, MPEF-PI 6132.18, Biozone NP4, upper Danian, Roca Formation, General Roca, Río Negro (all scale bars: 300 μ m). **A**, general view of the colony. **B**, detail of the edge of the colony. **C**, autozooids with prominent median keels. **D–E**, two gonozooids from the same colony, with the ooeciopore arrowed in **E**.

originally subcircular. Gonozooids with conspicuously inflated brood chamber (Fig. 6D–E), two times wider than long, oval to rounded triangular with straight distal edge; roof penetrated by numerous autozooidal peristomes; ooeciopore located centrally, slightly smaller than an autozooidal aperture and somewhat transversely elongate. Narrow budding zone exposing one or two generations of autozooids. Protoecium and ancestrula not seen. Autozooids arranged in a roughly quincuncial pattern; frontal walls elongate, becoming more convex distally, zooidal boundaries generally distinct; pseudopores conspicuous (Fig. 6F), longitudinally oval; apertures subcircular, isolated; preserved peristomes short, inclined distally.

Measurements. AS $285 \pm 35 \mu\text{m}$ (1, 11); AD $58 \pm 7 \mu\text{m}$ (1, 10); BCL $750 \mu\text{m}$ (1, 1); BCW $1475 \pm 177 \mu\text{m}$ (1, 2).

Remarks. The transversely elongate brood chamber pierced by autozooids is typical of the genus *Plagioecia* among cyclostomes of the 'Berenicea' type (Taylor and Sequeiros, 1982). We defer introducing a new name for this species because a large number of closely similar bereniciform species have been described from the Late Cretaceous to Danian but in very few cases are these well enough known (e.g. gonozooids are often not described) to be accurately compared with the Roca species. It is beyond the scope of the present paper to revise these species in order to establish that the Roca species is new.

***Plagioecia* aff. *cristata* Taylor and McKinney, 2006**

Fig. 7A–E

Plagioecia aff. 2006 *Plagioecia cristata*; Taylor and McKinney, p. 31, pl. 13.

Material. MPEF-PI 6132.18. Colony encrusting a valve of *Ostrea wilckensi*; upper Danian, Roca Formation, General Roca, Río Negro.
Description. Colony encrusting, multiserial, subcircular in outline. Budding zone moderately broad, exposing two to three generations of autozooidal buds. Peripheral subcolonies developed at budding zone, subcircular, partly overgrowing parent colony (Fig. 7A, upper right). Ancestrula not visible: earliest astogenetic stages fouled by a small oyster in the single specimen studied. Autozooids arranged quincuncially, about three or four times longer than wide (Fig. 7B); frontal wall convex with a prominent median keel (Fig. 7C), autozooidal boundaries generally distinct; pseudopores conspicuous, longitudinally oval; apertures circular or longitudinally elongate, isolated, never connate; peristomes inclined distally, preserved length short. Gonozooids with conspicuously inflated brood chambers (Fig. 7D–E), transversely elongate, at least twice wider than long, penetrated by autozooidal peristomes around the edge and occasionally more centrally. Ooeciopore terminal, smaller than an autozooidal aperture, transversely elongated, about half the diameter of an autozooidal aperture.

Measurements. AS $285 \pm 35 \mu\text{m}$ (1, 11); AD $58 \pm 7 \mu\text{m}$ (1,10); BCL $750 \mu\text{m}$ (1, 2); BCW $1475 \pm 177 \mu\text{m}$ (1, 2).

Remarks. This species shows a close similarity to *Plagioecia cristata* Taylor and McKinney, 2006 from the Maastrichtian of the southeastern USA. However, the gonozooids are broader in specimens from the Roca Formation. The Roca species is also similar to *Plagioecia parvipora* (Canu, 1922), '*P. formosa* (Canu, 1922) and '*Plagioecia*' *antanihodiensis* Di Martino, Martha and Taylor, 2018 from the Maastrichtian of Madagascar. *P. parvipora* differs from *P. aff. cristata* in having a budding zone exposing only 1–2 generations of autozooids and in the flat frontal walls of the autozooids. '*Plagioecia*' *formosa* is the most similar of the Madagascan species to *P. aff. cristata*, but has a central, circular ooeciopore. Finally, '*Plagioecia*'

antanihodiensis colonies have autozooids with convex frontal walls and small circular pseudopores.

Genus *Mesenteripora* de Blainville, 1830

Type species: *Mesenteripora michelini* de Blainville, 1830, Bathonian, Calvados, Normandy, France (see Walter, 1970).

***Mesenteripora* sp.**

Fig. 8A–E

Material. MPEF-PI 6132.28, encrusting base of a colony on a valve of *Pycnodonte* (*Phygraea*) *sarmientoi*; upper Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colony base encrusting, multiserial, subcircular in outline, somewhat lobate, with six broken erect bifoliate fronds arranged radially (Fig. 8A). Early astogenetic stages present, ancestrula overgrown (Fig. 8B). Autozooids elongate, boundaries marked by a narrow ridge; frontal walls becoming more convex distally; preserved peristomes short, inclined distally; apertures arranged quincuncially, circular to longitudinally elongate, isolated, never connate (Fig. 8C). Gonozooid with inflated brood chamber (Fig. 8D), slightly wider than long, oval to semicircular with straight distal edge, incorporating peristomial bases of some neighbouring autozooids (Fig. 8E); ooeciopore located on distal edge of brood chamber, minute, subcircular. Secondary nanozooids evident as autozooids capped by terminal diaphragms with a tiny central aperture on a short peristome (Fig. 7D–E)

Measurements. AS $240 \pm 32 \mu\text{m}$ (1, 13); AD $77 \pm 20 \mu\text{m}$ (1, 13), OD $56 \mu\text{m}$ (1, 1), NAD $60 \pm 4 \mu\text{m}$ (1, 3).

Remarks. *Mesenteripora* has erect colonies with bifoliate branches, either broad and frondose or less often narrow and palmate. Only the bases of the branches are preserved in the Roca specimen which resembles *Mesenteripora lirella* Taylor and McKinney, 2006 from the Campanian of Delaware, USA, but differs in having secondary nanozooids and autozooids lacking a prominent median keel on their frontal walls.

Genus *Actinopora* d'Orbigny, 1851

Type species: *Actinopora regularis* d'Orbigny, 1851 (= *Ceripora stellata* Koch & Dunker, 1837), Valanginian, Sainte-Croix, Switzerland.

***Actinopora robertsoniana* Canu, 1911**

Fig. 9A–E

1911 *Actinopora robertsoniana* Canu, p. 276, pl. 12, fig. 3.

Material. MPEF-PI 6132.17, colony encrusting a valve of *Ostrea wilckensi*; upper Danian, Roca Formation, General Roca, Río Negro.
Description. Colony encrusting, discoidal, unilaminar (Fig. 9A). Autozooidal apertures arranged in radial fascicles separated by areas of exterior wall containing circular pseudopores; fascicles initially uniserial, becoming bi- or triserial (Fig. 9B); new fascicles intercalated between existing fascicles (Fig. 9C). Gonozooid (Fig. 9D) large, transversely elongate, narrow, distal edge almost straight, interrupting fascicles; brood chamber pierced by a single autozooidal peristome in figured example (Fig. 9E); ooeciopore terminal, obliquely transversely elongate, slightly larger than an autozooidal aperture.

Measurements. ASW $130 \pm 29 \mu\text{m}$ (1, 22); Gap $245 \pm 53 \mu\text{m}$ (1, 25); BCW $2933 \mu\text{m}$ (1, 1); BCL $533 \mu\text{m}$ (1, 1); OD $63 \mu\text{m}$ (1, 1).

Remarks. Encrusting cyclostomes attributed to *Actinopora* are very common in the Valanginian–Maastrichtian of Europe (e.g., Gregory, 1909), and the genus has also been recorded living at the present-day (Canu and Bassler, 1929). Colonies are discoidal, often




Fig. 8. *Mesenteripora* sp., MPEF-PI 6132.28, NP4 Biozone, upper Danian, Roca Formation, Casa de Piedra, La Pampa. **A**, general view of the colony (scale bar: 1 mm). **B**, early astogeny (scale bar: 200 μm). **C**, broken base of an erect branch (scale bar: 400 μm). **D**, secondary nanozooids (scale bar: 100 μm). **E**, gonozooid (oocypore arrowed) surrounded mostly by secondary nanozooids (scale bar: 400 μm).

develop peripheral subcolonies (Taylor et al., 2018, fig. 4f), have autozooidal apertures arranged in radial fascicles beyond the zone of astogenetic change, and possess gonozooids with transversely elongate brood chambers. In the absence of a modern revision of *Actinopora*, the relationships between the Roca species, which was named *A. robertsoniana* by Canu (1911), and other nominal species are unclear.

Family incertae sedis

'Berenicea' sp.

Fig. 9F

Material. MPEF-PI 6132.13, colony encrusting a valve of *Ostrea wilckensi*; upper Danian, Roca Formation, General Roca, Río Negro. **Measurements.** ZL $143 \pm 40 \mu\text{m}$ (1, 5); ZW $90 \pm 10 \mu\text{m}$ (1, 5).

Description. Colony encrusting, multiserial, unilaminar, immature, fan-shaped, measuring up to 500 μm in diameter, infertile; two generations of buds visible at growing edge. Autozooids about 100 μm long by 90 μm wide (Fig. 9F). Frontal wall convex. Peristomes broken. Ancestrula preserved, not overgrown, protoecium 100 μm in width.

Remarks. The species is left in open nomenclature, following the recommendations of Taylor and Sequeiros (1982) for infertile tubuliporines with 'bereniciform' colonies which, depending on the morphology of the gonozooid, may belong to *Microeciella* Taylor and Sequeiros, 1982, *Reptomultisparsa* d'Orbigny, 1851, *Hyporosopora* Canu and Bassler 1929, *Mesonopora* Canu and Bassler, 1929 or *Plagioecia*.

Suborder Rectangulata Waters, 1887

Family Lichenoporidae Smitt, 1866

Genus *Disporella* d'Orbigny, 1851

Type species: *Discopora hispida* Fleming, 1828, Recent, British Isles.

?*Disporella discoidea* (Canu, 1911)

Fig. 9G–H

1911 *Reptocavea discoidea* Canu, p. 278, pl. 12, figs 1–2.

Material. MPEF-PI 6132.65, colony encrusting a valve of *Pycnodonte (Phygraea) sarmientoi*; upper Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colony encrusting, macula at centre slightly depressed (Fig. 9G). Autozooidal apertures longitudinally oval, arranged quincuncially. Alveoli small, located between autozooids and filling macula (Fig. 9H). Gonozooid not seen.

Measurements. OL $82 \pm 7 \mu\text{m}$ (1, 7); OW $52 \pm 4 \mu\text{m}$ (1, 7).

Remarks. The taxonomy of rectangulate cyclostomes is difficult and generic definitions are vague. Therefore, the assignment of Canu's (1911) species to *Disporella* is very tentative, made more so by the lack of a gonozooid.

Suborder Cerioporina von Hagenow, 1851

Family Cerioporidae Busk, 1859

Genus *Ceriopora* Goldfuss, 1826

Type species: *Ceriopora micropora* Goldfuss, 1826, Maastrichtian, Maastricht, Netherlands (see Nye, 1976).

Ceriopora sp.

Fig. 10A–C

Material. MPEF-PI 6132.2, colony encrusting a valve of *Cubitostrea ameghinoi*; upper Danian, Roca Formation, Bajada de Jagüel, Neuquén.

Description. Colony encrusting, massive, compound, multilayered (Fig. 10A). Zooidal apertures polygonal, variable in size, lacking any clear dimorphism between autozooids and kenozooids (Fig. 10B–C). Gonozooid not seen.

Measurements. AS $109 \pm 13 \mu\text{m}$ (1, 35); AD $76 \pm 20 \mu\text{m}$ (1, 35).

Remarks. The genus *Ceriopora* has been used in a broad sense to include erect and massive encrusting species in which a clear distinction between larger autozooids and smaller kenozooids is typically lacking, as in the species described here from the Roca Fm. The type species of *Ceriopora*, *Ceriopora micropora*, was redescribed by Nye (1976), with emphasis on internal morphological characters (Hara, 2001).

Order Cheilostomata Busk, 1852

Suborder Membraniporina Ortmann, 1890

Superfamily Membraniporoidea Busk, 1854

Family Electridae Stach, 1937

Genus *Electra* Lamouroux, 1816

Type species: *Flustra verticillata* Ellis and Solander, 1786, Recent, North Atlantic Ocean (see Nikulina et al., 2012).

Electra sp.

Fig. 10D–E

Material. MPEF-PI 6132.1, colony encrusting a valve of *Cubitostrea ameghinoi*; upper Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, forming irregular patches on the substrate. Autozooids elongate pyriform in frontal outline shape (Fig. 10D). Gymnocyst well-developed, in some zooids bearing a short spine on the median proximal border of the opesia (Fig. 10E). Cryptocyst narrow. Opesia longitudinally elliptical. **Measurements.** ZL $314 \pm 22 \mu\text{m}$ (1, 9); ZW $200.00 \pm 22 \mu\text{m}$ (1, 11); OpL $210 \pm 24 \mu\text{m}$ (1, 10); OpW $163 \pm 13 \mu\text{m}$ (1, 7).

Remarks. The studied specimen resembles *Electra everretti* Taylor and McKinney, 2006 from the Maastrichtian of North Carolina, USA. However, *E. everretti* has a wider cryptocyst and more elongate autozooids. The fragmentary colony from Roca Fm. consists of only 18 to 20 poorly preserved zooids. While the Roca Fm. species, along with *E. everretti*, may be closer to *Einhornia* Nikulina, 2007, the original diagnosis of *Einhornia* stipulates the presence of calcified opercula, for which there is no evidence. It is therefore assigned to *Electra* in the broad sense of the genus. Nevertheless, the presence of a spine on the medioproximal edge of the opesia of some of these zooids allows it to be assigned to *Electra sensu lato*.

Genus *Conopeum* Gray, 1848

Type species: *Millepora reticulum* Linnaeus, 1767, Recent, North Atlantic Ocean.

Conopeum okaiana (Canu, 1911)

Fig. 10F–H

1911 *Membranipora okaiana* Canu, p. 224, pl. 2, fig. 10.

Material. GHUNLPam 17423, colony encrusting a valve of *Turkostrea argentina* Griffin et al., 2005; lower Danian, Roca Formation, Liu Malal, Mendoza.

Description. Colony encrusting, multiserial, unilamellar (Fig. 10F). Autozooids arranged in well-defined longitudinal rows; broad and rounded hexagonal in frontal outline shape, usually longer than




Fig. 9. A–E. *Actinopora robertsoniana* Canu, 1911, MPEF-PI 6132.17, NP4 Biozone, upper Danian, Roca Formation, General Roca, Río Negro. A, general view of colony (scale bar: 1 mm). B, detail (scale bar: 40 μ m). C, fascicle and growing edge (scale bar: 200 μ m). D, gonozooid of a second colony (scale bar: 400 μ m). E, detail showing probable oocypore (arrowed) (scale bar: 200 μ m). F. '*Berenicea*' sp., MPEF-PI 6132.13, NP4 Biozone, upper Danian, Roca Formation, General Roca, Río Negro (scale bar: 200 μ m). G–H. *Disporella? discoidea* (Canu, 1911), MPEF-PI 6132.65, NP4 Biozone, upper Danian, Roca Formation, Casa de Piedra, La Pampa. G, general view (scale bar: 400 μ m). H, detail showing kenozooids (alveoli) separating autozooidal apertures (scale bar: 100 μ m).

wide; gymnocyst lacking; cryptocyst broadest proximally, narrowing distally, not forming a distinct shelf; opesia oval or inverted pear-shaped, mural rim crenulated; imperforate closure plates present in some autozooids (Fig. 10H). Ancestrula and early astogeny unknown.

Measurements. ZL $235 \pm 88 \mu\text{m}$ (1, 28); ZW $175 \pm 67 \mu\text{m}$ (1, 27); OpL $302 \pm 21 \mu\text{m}$ (1, 6); OpW $205 \pm 32 \mu\text{m}$ (1, 6).

Remarks. *Conopeum spissamentum* Taylor and McKinney, 2006, from the Late Cretaceous of the southeastern USA, differs from *Conopeum okaiana* in normally having small kenozooids at the proximolateral corners of the autozooids which are more rectangular in outline shape. The closure plates of *C. spissamentum* and of two other species from the Late Cretaceous of the southeastern USA – *Conopeum nelsoni* Canu and Bassler, 1926 and *Conopeum paranelsoni* Taylor and McKinney, 2006 – are porous, unlike those of *C. okaiana*.

Suborder Flustrina Smitt, 1868

Superfamily Calloporoidea Norman, 1903

Family Calloporidae Norman, 1903

Genus *Flustrellaria* d'Orbigny, 1851

Type species: *Flustrellaria fragilis* d'Orbigny, 1851, Cenomanian. Le Mans, France.

***Flustrellaria* sp.**

Fig. 11A–C

Material. MPEF-PI 6132.21, colony encrusting a valve of *Pycnodonte (Phygraea) vesicularis* Lamarck, 1806; upper Maastrichtian, Jagüel Formation, Bajada del Jagüel, Neuquén.

Description. Colony encrusting, multiserial, unilaminar. Autozooids rounded rhomboidal in frontal outline, arranged more-or-less quincuncially (Fig. 11A); gymnocyst poorly developed; spine bases numbering about 8–10 (Fig. 11B), distributed circumopesially; cryptocyst finely pustulose, narrowing slightly distally; opesia longitudinally elliptical to pear-shaped. Ovicells broken and crushed (Fig. 10B). Interzooidal avicularia about half the size of autozooids, oval or inverted pear-shaped, rostrum apparently short and rounded (Fig. 11C). Ancestrula and early astogeny not observed. **Measurements.** ZL $395 \pm 26 \mu\text{m}$ (1, 10); ZW $263 \pm 21 \mu\text{m}$ (1, 10); AL $327 \mu\text{m}$ (1, 1); AW $145 \mu\text{m}$ (1, 1); OpL $250 \pm 26 \mu\text{m}$ (1, 10); OpW $165 \pm 24 \mu\text{m}$, 15 (1, 10).

Remarks. *Flustrellaria* sp. differs from the US Late Cretaceous species *F. anatina* Canu and Bassler, 1926, which has large, spatulate interzooidal avicularia. However, the number of spines can be the same in both species (8–10), although these often include a larger mid distolateral pair in *F. anatina* (see Taylor and McKinney, 2006, pl. 50, fig. 1C).

Genus *Pyriporella* Canu, 1911

Type species: *Pyriporella ameghinoi* Canu, 1911, Danian, Argentina.

***Pyriporella ameghinoi* Canu, 1911**

Fig. 11D–G

1911 *Pyriporella ameghinoi* Canu, p. 235, pl. 4, figs 8–9.

1911 *Pyriporella confluens* Canu, p. 236, pl. 4, figs 10–11.

Material. MPEF-PI 6132.18, colony encrusting a valve of *Ostrea wilckensi*; upper Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, unilaminar. Autozooids arranged more-or-less quincuncially, elongate pear-shaped

(Fig. 11D); gymnocyst moderately developed proximally, narrowing laterally and absent distally, spine bases lacking; cryptocyst broad, proximally decreasing in width distally, sloping gently inwards; opesia oval, occupying approximately half of frontal surface area. Ancestrula subcircular, budding a distal and two distolateral zooids, no or few avicularia in early astogeny (Fig. 11E). Intramural buds present in a few zooids (Fig. 11F). Closure plates not seen. Ovicell hyperstomial, large, ectoecium usually completely calcified, a median fissure often evident (Fig. 11G). Avicularia interzooidal, small, numerous, located at corners of autozooids, oriented distolaterally, rounded, pivotal bar not calcified (Fig. 11G).

Measurements. ZL $329 \pm 88 \mu\text{m}$ (1, 15); ZW $257 \pm 98 \mu\text{m}$ (1, 15); AL $140 \pm 140 \mu\text{m}$ (1, 12); AW $70 \pm 18 \mu\text{m}$ (1, 12); OvL $206 \pm 14 \mu\text{m}$ (1, 6); OvW $253 \pm 18 \mu\text{m}$ (1, 5).

Remarks. The paratype of *Pyriporella confluens* Canu, 1911 (MACN-Pi 1844) has the same features as *P. ameghinoi* and is considered to be a junior synonym.

Family Antroporidae Vigneaux, 1949

Genus *Akatopora* Davis, 1934

Type species: *Akatopora clausentina* Davis, 1934, Eocene, Hampshire, England (see Gordon 1986, plate 7E).

?***Akatopora* sp. 1**

Fig. 12A–B

Material. MPEF-PI 6132.20, colony on a valve of *Amphidonte mendozana* Ihering, 1907; upper Maastrichtian, Roca Formation, Huantraico, Neuquén.

Description. Colony encrusting, multiserial, unilaminar (Fig. 12A). Autozooids quincuncially arranged and surrounded by smaller polymorphs that overlap their edges; longitudinally elongate; gymnocyst and cryptocyst not visible due to poor preservation and presence of polymorphs; opesia occupying most of frontal surface, oval. Ovicells not observed. Polymorphs, presumed to be avicularia, numbering about 2–4 per autozooid, directed distally or laterally (Fig. 12B).

Measurements. OpL $325 \pm 34 \mu\text{m}$ (1, 6); OpW $15 \pm 22 \mu\text{m}$ (1, 6); AL $77 \pm 11 \mu\text{m}$ (1, 11); AW $61 \pm 13 \mu\text{m}$ (1, 11).

Remarks. This specimen shows similarities with *Akatopora granulata* (Canu, 1911), but the poor preservation does not allow either the generic or specific identity to be confirmed.

***Akatopora kaufmanni* sp. nov.**

Fig. 12C–E

Etymology. Named for the collector of the holotype, W. Kaufmann. **Diagnosis.** *Akatopora* with very small autozooids; cryptocyst moderately wide, granular; opesia pear shaped.

Type horizon. Roca Fm. (Paleocene, upper Danian).

Material. Holotype: NHMUK D32352a, Danian, Roca Formation, General Roca, Río Negro; collected by W. Kaufmann and purchased May 1927.

Description. Colony encrusting, multiserial, unilaminar (Fig. 12C). Autozooids surrounded by smaller polymorphs that overlap their edges (Fig. 12D); longitudinally elongate; gymnocyst reduced, broadest proximally and tapering distally along the margins of the zooids; cryptocyst moderately wide, tapering distally, inwardly sloping, granular; opesia occupying most of frontal surface, oval or pear-shaped. Intramural buds present but closure plates not observed. Ovicells hyperstomial, ectoecium fully calcified,




Fig. 10. A–C. *Ceripora* sp., MPEF-PI 6132.2, NP4 Biozone, upper Danian, Roca Formation, Bajada de Jagüel, Neuquén. **A**, general view of the colony (scale bar: 1 mm). **B**, undifferentiated autozooids and kenozooids (scale bar: 200 μ m). **C**, detail of zooidal apertures (scale bar: 100 μ m). D–E. *Electra* sp., MPEF-PI 6132.1, NP4 Biozone, upper Danian, Roca Formation, General Roca, Río Negro. **D**, general view. **E**, group of autozooids with gymnocystal frontal walls. (scale bars: 200 μ m). F–H. *Conopeum okaiana* Canu, 1911, GHUNLPam 17423, NP1–NP2 Biozone, lower Danian, Roca Formation, Liu Malal, Mendoza. **F**, autozooids (scale bar: 400 μ m). **G**, hexagonal zooids (scale bar: 200 μ m). **H**, autozooids, one with a closure plate (scale bar: 200 μ m).

sometimes with a medial suture, wider than long, *c.* 70 μ m long by 110 μ m wide. Polymorphic zooids, which may be a mixture of avicularia and kenozooids, infilling most of the spaces between the autozooidal opesiae, variable in orientation and size, typically 80–130 μ m long by 50–90 μ m wide (Fig. 12E).

Measurements. ZL 284 \pm 44 μ m (1, 7); ZW 160 \pm 27 μ m (1, 7); OpL 187 \pm 8 μ m (1, 7); OpW 121 \pm 18 μ m (1, 7).

Remarks. This new species differs from both the US Maastrichtian species *Akatopora sulcata* (Canu and Bassler, 1926) (see Taylor and McKinney 2006, p. 88) and the New Zealand Recent species *Akatopora circumsaepa* (Uttley, 1951) (see Gordon 1986, p. 35) in its smaller autozooids with proportionally wider cryptocysts. Compared with the Maastrichtian species ?*Akatopora* sp. 1 described above, this Danian species also has smaller zooids. Together with *Cianotremella gigantea* Canu 1911, the holotype encrusts a Roca Formation oyster shell that was purchased by the NHMUK in 1927. Because the two species are so well preserved, they are described here even though the exact locality and stratigraphical horizon from which they were collected is uncertain.

Superfamily Microporoidea Gray, 1848

Family incertae sedis

Genus *Cianotremella* Canu, 1911

Type species: *Cianotremella gigantea* Canu, 1911, Danian, Roca Formation, Argentina.

***Cianotremella gigantea* Canu, 1911**

Fig. 12F–H

1911 *Cianotremella gigantea* Canu, p. 257, pl. 7, fig. 14.

Material. NHMUK D32352b, Danian, Roca Formation, General Roca, Río Negro; collected by W. Kaufmann and purchased May 1927.

Description. Colony encrusting, multiserial, unilaminar (Fig. 12F). Autozooids rectangular to elongate rhomboidal in shape, separated by thin, raised boundary walls; gymnocyst lacking; frontal wall an extensive cryptocyst occupying most of the frontal surface, convex, coarsely granular, apparently imperforate; opesia semicircular, proximal edge slightly bowed, a narrow distal oral shelf visible in some zooids; oral spines lacking. Ovicells absent in the studied specimen. One example of a vicarious kenozooid observed, a little narrower than the autozooids but about the same length; opesia longitudinally elliptical, slightly more than half of the length of the zooid; no associated avicularium. Avicularia present distally of most, possibly all, autozooids, transversely oriented, located within the boundary wall of the associated autozooid (Fig. 12G); rostrum with a rounded or subrounded tip, slightly curved to parallel the distal margin of the autozooidal opesia; a constriction dividing the avicularian opesia from the rostrum but no complete pivotal bars observed (Fig. 12H).

Measurements. ZL 550 \pm 62 μ m (1, 10); ZW 370 \pm 44 μ m (1, 10); OpL 123 \pm 8 μ m (1, 10); OpW 178 \pm 14 μ m (1, 10).

Remarks. Although *Cianotremella* is a distinctive genus, its family-level classification is uncertain. When introducing this monospecific genus, Canu (1911) believed it to be an ascophoran cheilostome. This opinion was followed by Bassler (1953) in the bryozoan *Treatise*. However, *Cianotremella* is clearly an anascan-grade cheilostome with an extensive cryptocystal frontal wall. It may be closely related to *Stictostega* (see Taylor and McKinney, 2006), with which it shares the presence of a small avicularium

distal to the autozooid, although the perforations seen in the frontal walls of *Stictostega* Shaw, 1967 cannot be observed in *Cianotremella*. The specimen described here appears to be infertile. Canu (1911, p. 258) described the ovicell thus: “Ovicelle cachée dans la partie supérieure de la zoécie, s’ouvrant, par une fente saillante et transverse, au-dessus de l’aperture”; i.e., immersed and opening via a transverse slit.

Family Aspidostomatidae Jullien, 1888

Genus *Aspidostoma* Hincks, 1881

Type species: *Aspidostoma crassum* Hincks, 1881 = *Eschara gigantea* Busk, 1854, Recent, South Atlantic.

***Aspidostoma onychocelliferum* Canu, 1911**

Fig. 13A–D

1911 *Aspidostoma onychocelliferum* Canu, p. 254, pl. 6, fig. 12.

Material. MPEF-PI 6132.68, encrusting colony on a valve of *Pycnodonte* (*Phygraea*) *burckhardti* (Böhm, 1903); lower Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, unilaminar. Autozooids rounded hexagonal, separated by deep grooves (Fig. 13A); cryptocystal frontal wall slightly convex, distally sloping inwards towards the opesia, granular (Fig. 13B); opesia wider than long, semi-elliptical. Ovicells hyperstomial, with a cryptocyst-like, non-porous surface. Ancestrula surrounded by six zooids, including a distal and two distolateral zooids budded directly from the ancestrula (Fig. 13C). Interzooidal avicularia small, elliptical, oriented parallel to the edges of the autozooids, pivotal bar not calcified (Fig. 13D).

Measurements. ZL 435 \pm 39 (1, 9); ZW 324 \pm 37 (1, 9); OpL 87 \pm 11 (1, 9); OpW 148 \pm 28 (1, 9); OvL 93 \pm 10 (1, 2); OvW 129 \pm 20 (1, 2); AL 133 \pm 38 (1, 3); AW 83 \pm 7 (1, 3).

Remarks. *Aspidostoma* is endemic to the Southern Hemisphere; Cretaceous species assigned to this genus from France belong elsewhere (Taylor, 2019). *Aspidostoma onychocelliferum* Canu, 1911 resembles the Eocene species, *Aspidostoma pyriformis* Hara, 2001, from the La Meseta Formation of the Antarctic Peninsula, but the autozooids of *A. pyriformis* are smaller, the opesiae have anvil-shaped processes, and the interzooidal avicularia are pyriform.

Superfamily Monoporelloidea Hincks, 1882

Family Monoporellidae Hincks, 1882

Genus *Monoporella* Hincks, 1881

Type species: *Haploporella nodulifera* Hincks, 1881, Recent, SE Australia (Cook et al., 2018, p. 129).

***Monoporella convexa* (Canu, 1911)**

Fig. 13E–H

1911 *Micropora convexa* Canu, p. 250, pl. 7, figs 1–3.

Material. MPEF-PI 6132.15 and MPEF-PI 6132.16, colonies encrusting valves of *Ostrea wilckensi*; upper Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting multiserial, unilaminar. Autozooids subhexagonal (Fig. 13E); opesia semielliptical, small, with low rim, lacking oral spine bases in zooids from zone of astogenetic repetition; frontal wall cryptocystal, slightly convex, finely granulated, penetrated evenly by about 8–10 opesiules. Ancestrula similar to later budded zooids but smaller and with four oral spine bases




Fig. 11. A–C. *Flustrellaria* sp., MPEF-PI 6132.21, upper Maastrichtian, Jagüel Formation, Bajada del Jagüel, Neuquén. A, general view (scale bar: 400 μ m). B, detail of zooids with small circumopesia spine bases and damaged ovicells (scale bar: 200 μ m). C, interzooidal avicularium (lower centre) (scale bar: 200 μ m). D–G. *Pyriporella ameghinoi* Canu, 1911, MPEF-PI 6132.18, NP4 Biozone, upper Danian, Roca Formation, General Roca, Río Negro. D, general view of the colony (scale bar: 1 mm). E, early astogeny, the ancestrula and first buds missing (scale bar: 100 μ m). F, autozooids, most ovicellate (scale bar: 200 μ m). G, ovicellate autozooids and small avicularia (scale bar: 200 μ m).

(Fig. 13F). Ovicells broken, the spinose roofs missing exposing the gymnocrystal floor (Fig. 13G–H).

Measurements. ZL $504 \pm 166 \mu\text{m}$ (3, 42); ZW $310.5 \pm 83 \mu\text{m}$ (3, 38); OpL $76 \pm 13 \mu\text{m}$ (3, 63); OpW $98 \pm 15 \mu\text{m}$ (3, 62).

Remarks. The observed characters of our material match those of the paratype of *M. convexa* (MACN-Pi 1859). *Monoporella chubuti* (Canu, 1911), also from the Roca Formation, has larger opesia. Revision of both *M. convexa* and *M. chubuti*, including scanning electron microscopy of type material, will be needed to calify the identity of these two species.

Superfamily ?Thalamoporelloidea Levinsen, 1902

Family ?Steginoporellidae Hincks, 1884

Genus ?*Labioporella* Harmer, 1926

Type species: *Labiopora crenulata* Levinsen, 1909, Recent, ?Torres Strait.

?*Labioporella* sp.

Fig. 14A–D

Material. MPEF-PI 6132.8, encrusting colony on a valve of *Pycnodonte* (*Phygraea*) *sarmientoi*; upper Danian, Formación Roca, Casa de Piedra, La Pampa.

Description. Colony encrusting, multiserial, unilaminar (Fig. 14A). Autozooids arranged in well-defined rows (Fig. 14B), rounded rectangular in outline shape, proximal edge concave, distal edge convex, variable in width (Fig. 14C), separated by grooves; gymnocrystal lacking; cryptocyst well developed, shelf-like, occupying about half of the length of the zooid, granular, sloping inwards at its distal end, narrow proximally; opesia semi-elliptical, longer than wide (Fig. 14D). Ovicells, avicularia and early astogeny not observed.

Measurements. ZL 259 ± 33 (1, 21); ZW 187 ± 27 (1, 25); OL 79 ± 10 (1, 23); OW 124 ± 33 (1, 23).

Remarks. This species is referred to *Labioporella* very tentatively. Although the form and arrangement of the autozooids, their granular cryptocystal frontal walls and opesiae that are longer than wide matches *Labioporella*, the species from the Roca Fm. lacks a prominent median process along the proximal edge of the opesia, and there is no indication of a polypide tube or tiny perforations in the cryptocyst, although these absences may be due to preservation. Should better evidence be found for placing the Roca species in *Labioporella*, it would represent the oldest known example of this genus, which at the present-day occurs in the Pacific, Indian and Atlantic oceans.

Superfamily Cribrilinoidea Hincks, 1879

Family Cribrilinidae Hincks, 1879

Genus *Tricephalopora* Lang, 1916

Type species: *Cribrilina triceps* Marsson, 1887, Maastrichtian, Rügen, Germany.

Tricephalopora sp. 1

Fig. 15A–C

Material. MPEF-PI 6132.19, colony encrusting a valve of *Gryphaeostrea callophyla* Ihering, 1903; lower Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, sheet-like. Autozooids elongate hexagonal; orifice round, gymnocrystal restricted to narrow band around perimeter of autozooid (Fig. 14A); frontal shield with

small costal field comprising 6–7 costae meeting and fusing along the midline of the zooid (Fig. 14B), no lateral intercostal fusions or pelmata visible; worn costae reveal the lumen. Adventitious avicularia paired, located at proximolateral official margin, directed proximolaterally inwards (Fig. 14C); rostrum triangular, blunt tipped; crossbar calcified. Ovicell smoothly inflated, lacking porous. Ancestrula and kenozooids not observed.

Measurements. ZL $447 \pm 41 \mu\text{m}$ (1, 13); ZW $298 \pm 26 \mu\text{m}$ (1, 13); OL $129 \pm 10 \mu\text{m}$ (1, 14); OW $139 \pm 12 \mu\text{m}$ (1, 17); AL $46 \pm 12 \mu\text{m}$ (1, 17); AW $59 \pm 9 \mu\text{m}$ (1, 17).

Remarks. This specimen is similar to *T. vibraculata* Turner, 1979 from the Late Cretaceous of New Jersey, USA. However, the specimen from Roca Fm. has fewer costae forming the frontal shield of the autozooids and lacks columnar ?kenozooids. The corroded costae mean that pelmata cannot be observed, and argues against introducing a new species for the Roca material.

?*Tricephalopora* sp.

Fig. 15D–E

Material. MPEF-PI 6132.3, colony encrusting a valve of *Cubitostrea ameghinoi*; upper Danian, Roca Formation, Cerros Bayos, La Pampa.

Description. Colony encrusting, multiserial, sheet-like. Autozooids elongate hexagonal (Fig. 14D); frontal shield with 12–18 costae meeting in a line along the midline of the zooid, no lateral intercostal fusions or pelmata. Scattered avicularia in various positions, poorly preserved. Ovicells gently inflated, lacking pores (Fig. 14E). Ancestrula not observed.

Measurements. ZL 545 ± 42 (1, 13) μm ; ZW $223 \pm 29 \mu\text{m}$ (1, 24); OL $137 \pm 11 \mu\text{m}$ (1, 13); OW $215 \pm 23 \mu\text{m}$ (1, 13); OvL $169 \pm 16 \mu\text{m}$ (1, 13); OvW $222 \pm 23 \mu\text{m}$ (1, 11).

Remarks. Specimens are poorly preserved, and typical features of the genus, such as paired avicularia on each autozooidal orifice seem to be lacking, hence assignment to *Tricephalopora* is tentative, even though this seems to be the most appropriate genus.

Infraorder Umbonulomorpha Gordon, 1989

Superfamily Arachnopusioidea Jullien, 1888

Family Arachnopusiidae Jullien, 1888

Genus *Poricella* Canu, 1904

Type species: *Poricella macconnica* Canu, 1904, Eocene, Tunisia.

Poricella tripora (Canu, 1911)

Fig. 16A–C

1911 *Hiantopora tripora* Canu, p. 256, pl. 7, fig. 4.

Material. MPEF-PI 6132.67, colony encrusting a valve of *Pycnodonte* (*Phygraea*) *sarmientoi*; upper Danian, Roca Formation, Bajada del Jagüel, Neuquén.

Description. Colony encrusting, multiserial, unilaminar. Autozooids elongate oval (Fig. 16A); frontal shield slightly convex (Fig. 16B), perforated by 3 foramina, often reniform due to the presence of a tongue-like process (Fig. 16C); orifice D-shaped, with a low peristome. Ovicells poorly preserved, apparently semicircular, wider than long. Adventitious avicularia present.

Measurements. ZL $384 \pm 34 \mu\text{m}$ (1, 16); ZW $226 \pm 28 \mu\text{m}$ (1, 15); OL $101 \pm 20 \mu\text{m}$ (1, 21); OW $137 \pm 19 \mu\text{m}$ (1, 21).

Remarks. This specimen has been compared with the paratype of *P. tripora* (MACN-Pi 1845) described by Canu (1911) with which it is deemed conspecific. Along with the following species, it may represent the oldest known record of the genus.




Fig. 12. A–B ?*Akatopora* sp. 1, MPEF-PI 6132.20, upper Maastrichtian, Roca Formation, Huantraico, Neuquén. **A**, general view of the colony (scale bar: 600 μ m). **B**, avicularia (scale bar: 200 μ m). C–E. *Akatopora kaufmanni* sp. nov., NHMUK D32352a. Danian, Roca Formation, General Roca, Río Negro. **C**, general view of the colony with an overgrowing colony of *Cianotremella gigantea* to the right (scale bar: 600 μ m). **D**, autozooids surrounded by small polymorphs (scale bar: 200 μ m). **E**, detail showing some autozooids with ovicells (200 μ m). F–H. *Cianotremella gigantea* Canu, 1911, NHMUK D32352b, Danian, Roca Formation, General Roca, Río Negro. **F**, general view of the colony fouled by a small *Poricella* and a worn spirorbid (lower right) (scale bar: 600 μ m). **G**, autozooids and avicularia (scale bar: 300 μ m). **H**, autozooids, avicularia and kenozooid (middle left) (scale bar: 400 μ m).

Poricella sp.

Fig. 16D–G

Material. MPEF-PI 6132.23, colony encrusting a valve of *Pycnodonte (Phygraea) sarmientoi*. Upper Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colony encrusting, multiserial, unilaminar (Fig. 16D); large distal pore chamber visible at growing edge. Zooids oval, convex; frontal shield perforated by 6–9 foramina, each with a tongue of calcification extending into the opening (Fig. 16E). Orifice D-shaped, slightly elongated, sometimes with a peristome. Adventitious avicularia oval or rounded rectangular, oriented proximally or somewhat proximolaterally, rostrum rounded, crossbar uncalcified (Fig. 16F–G). Early astogenetic stages preserved but not the ancestrula (Fig. 16F).

Measurements. ZL $317 \pm 41 \mu\text{m}$ (1, 17); ZW $223 \pm 34 \mu\text{m}$ (1, 18); OL $67 \pm 15 \mu\text{m}$ (1, 17); OW $70 \pm 12 \mu\text{m}$ (1, 17); AL $78 \pm 18 \mu\text{m}$ (1, 16); AW $80 \pm 18 \mu\text{m}$ (1, 16).

Remarks. Canu (1911, pl. 7, figs 12–13) illustrated material assigned to *Tremogasterina problematica*, which is similar to *Poricella* sp. According to curators at MACN, and as personally ascertained by one of us (S.S. Brezina), the type material of *T. problematica* is missing from the collection.

Genus *Tremogasterina* Canu, 1911

Type species: *Tremogasterina problematica* Canu, 1911. Danian, Roca Formation, General Roca, Argentina.

Tremogasterina problematica Canu, 1911

Fig. 17A–C

1911 *Tremogasterina problematica* Canu, p. 256, pl. 7, figs 12–13.
?1977 *Tremogasterina problematica* Canu, Cook, p. 127, pl. 3A–B.

Material. MPEF-PI 6132.11, colony encrusting a valve of *Pycnodonte (Phygraea) sarmientoi*; upper Danian, Roca Formation, Casa de Piedra, La Pampa.

Description. Colony encrusting, multiserial (Fig. 17A). Autozooids nearly equidimensional, frontal shield flat, pierced by 1–5 foramina (Fig. 17B–C), raised distally as a thickened bar proximal to the orifice. Orifice large, semicircular, rounded distally with an almost straight proximal edge. Ovicells subdued, ectoecium imperforate. Interzooidal avicularia numerous, oval (Fig. 17B), situated distolaterally of ovicells and laterally of autozooids, oriented distally or somewhat distolaterally, rostrum rounded, pivotal bar not calcified (Fig. 17D–E).

Measurements. ZL $281 \pm 30 \mu\text{m}$ (1, 4); ZW $219 \pm 22 \mu\text{m}$ (1, 4); OL $50 \pm 9 \mu\text{m}$ (1, 5); OW $78 \pm 10 \mu\text{m}$ (1, 5); AL $90 \mu\text{m} \pm 17 \mu\text{m}$ (1, 6); AW $94 \pm 19 \mu\text{m}$ (1, 6).

Remarks. Following the revision of Cook (1977), this is the only species now assigned to *Tremogasterina*, the others having been transferred to *Poricella*. The validity of *Tremogasterina* needs to be evaluated through a comparative study of *T. problematica* with the type species of *Poricella*, *P. macconnica* Canu, 1904, which is beyond the scope of the current study.

Genus *Trichinopolia* Guha and Nathan, 1996

Type species: *Trichinopolia crescentica* Guha and Nathan, 1996, Maastrichtian, Tamil Nadu, India.

?*Trichinopolia* sp.

Fig. 17F

Material. MPEF-PI 6132.20, colony encrusting a valve of *Amphidonte mendozana*; upper Maastrichtian, Roca Formation, Huantraico, Neuquén.

Description. Colony encrusting, multiserial. Autozooids hexagonal, slightly elongated, uniform in size and shape; frontal shield coarsely preserved, planar, apparently containing at least three foramina; orifice bell-shaped, the rim slightly raised. Adventitious avicularia paired laterally of the orifice which they indent slightly (Fig. 17F). Ovicells and ancestrula not observed.

Measurements. OL $94 \pm 11 \mu\text{m}$ (1, 8); OW $103 \pm 21 \mu\text{m}$ (1, 8); AL $80 \pm 10 \mu\text{m}$ (1, 7); AW $71 \pm 6 \mu\text{m}$ (1, 7).

Remarks. The studied specimen is of Maastrichtian age and resembles *Trichinopolia*, a genus first described from the Late Cretaceous of India and subsequently recorded from the Campanian–Maastrichtian of the southeastern USA and California (Taylor, 2008). In particular, the species from the Roca Fm. has an orifice of similar shape to *T. californica* Taylor, 2008 but seemingly lacks the prominent distal spine base seen in the North American species. Unfortunately, the preservation is too poor to be trustworthy about its generic identity. Nevertheless, *Trichinopolia* is the closest fit.

Superfamily Lepralielloidea Vigneaux, 1949

Family Romancheinidae Jullien, 1888

Genus *Balantiostoma* Marsson, 1887

Type species: *Cellepora marsupium* von Hagenow, 1839, Maastrichtian, Rügen, Germany.

Balantiostoma spectabilis (Canu, 1911)

Fig. 17G–H

1911 *Hoplocheilina spectabilis* Canu, p. 262, pl. 8, figs. 1–4.

Material. MPEF-PI 6132.15 and MPEF-PI 6132.66, colonies encrusting valves of *Ostrea wilkensi* Ihering, 1907; upper Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, unilaminar (Fig. 17G). Autozooids elongate hexagonal; frontal shield gently convex, granular, with large marginal areolar pores. Adventitious avicularia with rounded rostra, located singly or paired distolaterally of autozooidal orifice (Fig. 17H), pivotal bar calcified. Early astogeny and ovicells unknown.

Measurements. ZL $322 \pm 41 \mu\text{m}$ (2, 16); ZW $196 \pm 31 \mu\text{m}$ (2, 17); OL $94 \pm 31 \mu\text{m}$ (2, 13); OW $92 \pm 21 \mu\text{m}$ (2, 13); AL $36 \pm 0 \mu\text{m}$ (1, 4); AW $41 \pm 5 \mu\text{m}$ (1, 6).

Remarks. According to curators at MACN, and as personally ascertained by one of us (S.S. Brezina), the holotype of *Balantiostoma spectabilis* is unfortunately missing from the collection.

Balantiostoma elongata (Canu, 1911)

Fig. 18A–D

1911 *Exochella elongata* Canu, p. 264, pl. 9, figs. 1–3.

Material. MPEF-PI 6132.6, colony encrusting a valve of *Pycnodonte (Phygraea) sarmientoi*; upper Danian, Roca Formation, Casa de Piedra, La Pampa. MPEF-PI 6132.12, colony encrusting a valve of *Ostrea wilkensi*; upper Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, unilaminar. Early astogeny preserved; ancestrula ascophoran, small, circular, budding a distal and two distolateral zooids (Fig. 18A). Autozooids




Fig. 13. A–D. *Aspidostoma onychocelliferum* Canu, 1911., MPEF-PI 6132.68, lower Danian, Roca Formation, General Roca, Río Negro. A–B, general views of the colony (scale bars: A = 400 μ m; B = 200 μ m). C, early astogeny with first generations of zooids. D, ovicells and avicularia (scale bars: 200 μ m). E–H. *Monoporella convexa* Canu, 1911, MPEF-PI 6132.15, upper Danian, Roca Formation, General Roca, Río Negro. E, general view of autozooids (scale bar: 200 μ m). F, ancestrula (scale bar: 100 μ m). G, broken ovicell (scale bar: 100 μ m). H, early astogeny (scale bar: 200 μ m).

elongate, zooidal boundaries marked by grooves and areolar pores (Fig. 18B); frontal shield convex, primary orifice mucronate; without oral spines (Fig. 18B). Ovicell hyperstomial, small, globose, present in the majority of zooids (Fig. 18C). Adventitious avicularia sparse and scattered, located close to autozooidal orifices, directed laterally, small, transversely elliptical, pivotal bar not calcified (Fig. 18D).

Measurements. ZL $283 \pm 27 \mu\text{m}$ (2, 20); ZW $202 \pm 29 \mu\text{m}$ (2, 22); OL $101 \pm 11 \mu\text{m}$ (2, 16); OW $111 \pm 14 \mu\text{m}$ (2, 16); AL $39 \pm 48 \mu\text{m}$ (1, 2); AW $69 \pm 31 \mu\text{m}$ (1, 2); OvL $103 \pm 16 \mu\text{m}$ (1, 6); OvW $115 \pm 56 \mu\text{m}$ (1, 6).

Remarks. This species is similar to *Balantiostoma nomas* (Shaw, 1967) in having elongate autozooids delineated by prominent grooved boundaries and a mucronate orifice. However, it lacks the oral spines seen in this Late Cretaceous species from North America (see Taylor and McKinney, 2006).

Balantiostoma sp.

Fig. 18E–H

Material. MPEF-PI 6132.7, colony encrusting a valve of *Pycnodonte (Phygraea) sarmiento*; upper Danian, Roca Formation, Bajada del Jagüel, Neuquén. MPEF-PI 6132.13 and MPEF-PI 6132.14, colonies

encrusting valves of *Ostrea wilkensi*; Danian, Roca Formation, General Roca, Río Negro.

Description. Colony encrusting, multiserial, unilaminar (Fig. 18E). Early astogeny preserved, ancestrula tatiform with 78 spines, apparently budding a distal and two distolateral zooids (Fig. 18F). Autozooids elongate, rhomboidal; frontal shield granular; orifice usually mucronate; 4–6 oral spine bases, generally equal-sized (Fig. 18G–H). Ovicells only observed in MPEF-PI 6132.7, globose. Avicularia present in some autozooids, mostly those with ovicells (Fig. 18H); adventitious, located proximolaterally of opesia, unpaired, cystid chamber bulbous, rostral plane steeply inclined to colony surface.

Measurements. ZL $238 \pm 37 \mu\text{m}$ (3, 28); ZW $190 \pm 31 \mu\text{m}$ (3, 24); OL $54 \pm 19 \mu\text{m}$ (3, 23); OW $63 \pm 21 \mu\text{m}$ (3, 22); OvL $109 \pm 22 \mu\text{m}$ (1, 11); OvW $135 \pm 8 \mu\text{m}$ (1, 11); AL $111 \pm 16 \mu\text{m}$ (1, 13); AW $98 \pm 14 \mu\text{m}$ (1, 14).

Remarks. The studied specimens are moderately well-preserved and share some features with *Balantiostoma octospinigera* Taylor and McKinney, 2006 from the North American Cretaceous. However, the Roca Fm. species differs in having fewer oral spine bases (4–6 vs. 8) and a granular frontal shield.

Family incertae sedis




Fig. 14. ?*Labioporella* sp., MPEF-PI 6132.8, NP4 Biozone, Danian, Formación Roca, Casa de Piedra, La Pampa. A, general view of the colony (scale bar: 400 μ m). B–D, autozooids with semi-elliptical opesiae and shelf-like proximal cryptocysts arranged in longitudinal rows (scale bars: 200 μ m).




Fig. 15. A–C. *Tricephalopora* sp. 1. MPEF-PI 6132.19, lower Danian, Roca Formation, General Roca, Río Negro. **A**, general view of zooids (scale bar: 400 μ m). **B**, autozooids and adventitious avicularia located at proximolateral orificial margins (scale bar: 200 μ m). **C**, detail of costate frontal shield (scale bar: 100 μ m). **D–E**. ?*Tricephalopora* sp., MPEF-PI 6132.3, upper Danian, Roca Formation, Cerros Bayos, La Pampa. **D**, general view of the colony. **E**, ovicellate zooids (scale bars: 200 μ m).

Genus ***Eoporella*** gen. nov

Type species: *Eoporella lunata* sp. nov., upper Danian, Roca Formation, Casa de Piedra, La Pampa, Argentina.

Etymology. *Eos* (Gr.) for dawn, in reference to its similarity with the geologically older genus *Microporella*.

Diagnosis. As for type and only species.

Remarks. This new genus is introduced for a distinctive species from the Roca Fm. that strongly resembles the early Miocene–Recent genus *Microporella* but differs in several important morphological details and antedates the oldest known species of *Microporella* by about 40 million years. Condyles in *Eoporella* are situated distally of

the proximolateral corners, a little way along the edges of the orifice, whereas when present in *Microporella* they are in the proximolateral corners. The ovicell of *Eoporella* has a coarsely porous, smooth-surfaced entoecium separated from the frontal shield of the distal zooid by a narrow crescent of ectoecium. In contrast, the ovicell of *Microporella* comprises a granular, cryptocyst-like calcified wall that is continuous with the frontal shield of the distal zooid and can be imperforate or pierced by relatively small pores. Furthermore, most – though not all – species of *Microporella* have adventitious avicularia and oral spines, both lacking in *Eoporella*.




Fig. 16. A–C. *Poricella tripora* Canu, 1911, MPEF-PI 6132.67, upper Maastrichtian, Jagüel Formation, Bajada del Jagüel, Neuquén. **A**, general view of oval zooids, with broad orifice (scale bar: 200 μ m). **B**, adventitious avicularia (scale bar: 200 μ m). **C**, detail of frontal shield perforated by three foramina (scale bar: 60 μ m). **D–G**. *Poricella* sp., MPEF-PI 6132.23, upper Danian, Roca Formation, Casa de Piedra, La Pampa. **D**, general view of a colony (scale bar: 400 μ m). **E**, growth rim at the upper section. **F**, early astogenetic stages with small zooids (lower center). **G**, group of autozooids, some with peristomes, and avicularia. (scale bars: 200 μ m).




Fig. 17. A–E. *Tremogasterina problematica* Canu, 1911, MPEF-PI 6132.11, upper Danian, Roca Formation, Casa de Piedra, La Pampa. A, general view of the colony (scale bar: 400 μ m). B, oval interzooidal avicularia near the orifices (scale bar: 200 μ m). C, frontal shields with five foramina, highly abraded (scale bar: 100 μ m). D, zooids near growing edge, some with broken ovicells (scale bar: 200 μ m). E, growing edge with large distal pore chamber visible in upper centre (scale bar: 200 μ m). F, *?Trichinopolia* sp., MPEF-PI 6132.20, upper Maastrichtian, Roca Formation, Huantraico, Neuquén. Detail of frontal shield with poorly preserved lateroproximal adventitious avicularia (scale bar: 100 μ m). G–H. *Balantiostoma spectabilis* Canu, 1911, MPEF-PI 6132.15, upper Danian, Roca Formation, General Roca, Río Negro. G, general view of a small colony (scale bar: 400 μ m). H, detail of autozooids with areolae; avicularia proximolateral to the orifice (scale bar: 200 μ m).




Fig. 18. A–D. *Balantiostoma elongata* Canu, 1911, MPEF-PI 6132.6, upper Danian, Roca Formation, Casa de Piedra, La Pampa, upper Danian. A, early astogeny with putative ancestrula marked by an asterisk (scale bar: 200 μ m). B, detail of mucro in autozooidal orifice (scale bar: 100 μ m). C, elongate autozooids and areolar pores (scale bar: 200 μ m). D, small adventitious avicularia (scale bar: 100 μ m). E–H. *Balantiostoma* sp., MPEF-PI 6132.7, upper Danian, Roca Formation, Bajada del Jagüel, Neuquén. MPEF-PI 6132.13 and MPEF-PI 6132.14, upper Danian, Roca Formation, General Roca, Río Negro. E, general view of a colony (scale bar: 1 mm). F, early astogeny with ancestrula (marked by an asterisk) preserving long spines (scale bar: 100 μ m). G, zooids with mucro in proximal edge of the orifice (scale bar: 100 μ m). H, ovicellate zooids with avicularia proximolaterally of the orifice (scale bar: 200 μ m).




Fig. 19. *Eoporella lunata* gen. et sp. nov., MPEF-PI 6132.6, upper Danian, Roca Formation, Casa de Piedra, La Pampa. **A**, general view of a colony (scale bar: 600 μm). **B**, group of zooids (scale bar: 500 μm). **C**, zooids showing pseudoporous frontal shields (scale bar: 200 μm). **D**, orifice and crescent-shaped ascopore (scale bar: 200 μm). **E**, autozooids, some with broken ovicells (scale bar: 300 μm). **F**, ovicell with irregular pores in the entoecium (scale bar: 100 μm).

Eoporella lunata sp. nov.

Fig. 19 A–F

Etymology. Referring to the lunate shape of the ascopore.

Diagnosis. Lepralioid cheilostome with crescentic ascopore set in an extensive, non-porous concavity on the distal frontal shield bounded by a proximal continuation of the peristome. Orifice with lateral condyles. Ovicell hyperstomial, ectoecium uncalcified

except for a narrow rim, entoecium smooth with scattered large, irregular pores. Pore chambers lacking. No avicularia.

Type horizon. Roca Fm. (Paleocene, upper Danian).

Material. Holotype: MPEF-PI 6132.6; paratype: MPEF-PI 6132.10; encrusting colonies on valves of *Pycnodonte* (*Phygraea*) *sarmientoi*, upper Danian, Roca Formation, Casa de Piedra, La Pampa, Argentina.

Description. Colony encrusting, multiserial, unilaminar; pore chambers lacking. Autozooids rhomboidal; boundary walls well

raised (Fig. 19A–B); frontal shield convex, pseudoporous, pseudopores large, numbering about 40 per zooid; distinct areolar pores not discernible; orifice bell shaped, longer than wide, lateral condyles dividing an horseshoe-shaped anter from a broad, shallow sinus; no oral spines; peristome prominent and flared (Fig. 19C–D); ascopore large, crescentic, set in an extensive, non-porous, apron-like concavity on the frontal shield bounded by a proximal continuation of the peristome (Fig. 19D). Ovicell hyperstomial (Fig. 19E); ectoecium uncalcified except for a narrow, crescentic, smooth rim; entoecium smoothly calcified, with large but sparse (c. 10) pores irregularly oval in shape (Fig. 19F). Avicularia not observed.

Measurements. ZL $544 \pm 107 \mu\text{m}$ (2, 13); ZW $386 \pm 74 \mu\text{m}$ (2, 13); OL $103 \pm 39 \mu\text{m}$ (2, 15); OW $100 \pm 18 \mu\text{m}$ (2, 15); OpL $29 \pm 8 \mu\text{m}$ (2, 6); OpW $49 \pm 14 \mu\text{m}$ (2, 6); OvL $256 \pm 32 \mu\text{m}$ (2, 1); OvW $254 \pm 24 \mu\text{m}$ (2, 1); OOW $150 \mu\text{m}$ (1, 1).

Remarks. This is the only cheilostome in the Roca Fm. with a lepralioid frontal shield pierced by pseudopores. Indeed, *E. lunata* is believed to be the oldest lepraliomorph-grade cheilostome, which is the most speciose group of cheilostomes living in modern seas.

5. Conclusions

Our findings show that in northern Patagonia the diversity of encrusting bryozoans associated with oyster shells exhibits no major change across the K/Pg boundary (Table 1), in common with the pattern reported for northern Europe and the southeastern United States by Sogot et al. (2013, 2014). However, an important increase in the diversity is recorded during the upper Danian: at this time 13 cyclostome and 11 cheilostome species are documented on oyster shells contrasting with just four cheilostomes (i.e. *Flustrellaria* sp., *Akatopora* sp. 1, *Poricella tripora* and *Trichinopolia* sp.) on Maastrichtian oysters and three (i.e. *Conopeum okaiana*, *Aspidostoma onychocelliferum* and *Triccephalopora* sp. 1) on lower Danian shells. There are no records of Maastrichtian encrusting cyclostomes in the studied area, although the relationship between cyclostome and cheilostome species number in the Danian is congruent with information from the Northern Hemisphere (i.e. there are more species of cheilostomes after the K-Pg boundary). It is intriguing that cyclostomes slightly dominated over cheilostomes during upper Danian even though the record is limited. The lower relative abundance of cheilostomes during the upper Danian in the Neuquén Basin is coincident with the decline recorded at the end of the Danian in the Northern Hemisphere (Taylor and Waeschenbach, 2015).

The change in bryozoan biodiversity after the lower Danian correlates with an increase in the number of species of corals, molluscs, echinoids and crabs derived from low latitudes, reflecting higher seawater temperatures spreading south (Kiessling et al., 2005; Casadío et al., 2005; Aguirre-Urreta et al., 2008; Carrera and Casadío, 2016) coincident with sea surface temperatures obtained by Woelders et al. (2017) that show a warming phase after the K/Pg boundary in the Neuquén Basin.

Author statement

The authors S. Brezina, P. Taylor, M. Romero, E. Palópolo and S. Casadío, certify that all of us have seen and approved the final version of the manuscript, which is being submitted.

We warrant that the article is the authors' original work, has not received prior publication and is not under consideration for publication elsewhere.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by a PhD fellowship of the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) to S.S. Brezina. We also thank Claudia del Rio, who gave us access to the Canu Collection material from Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”. We greatly appreciate the corrections, suggestions and comments of the reviewers that significantly improved this work. Financial support was provided by PICT-2018-00917.

References

- Aguirre-Urreta, M.B., Casadío, S., Cichowolski, M., Lazo, D.G., Rodríguez, D., 2008. Afinidades paleobiogeográficas de los invertebrados cretácicos de la cuenca Neuquina. *Ameghiniana* 45, 591–611.
- Bassler, R.S., 1952. Taxonomic notes on genera of fossil and recent Bryozoa. *Journal of the Washington Academy of Sciences* 42, 381–385.
- Bassler, R.S., 1953. Bryozoa. In: Moore, R.C. (Ed.), *Treatise on invertebrate paleontology*. Part G. Geological Society of America and University of Kansas Press, Boulder and Lawrence, p. 253.
- de Blainville, H.M.D., 1830. Zoophytes. *Dictionnaire des Sciences Naturelle* 60, 1–631.
- Böhm, J., 1903. Über Ostreen von General Roca am Río Negro. *Zeitschrift der Deutsche geologische Gesellschaft* 55, 71–72.
- Brezina, S., Romero, M.V., Casadío, S., Bremec, C., 2014. Boring polychaetes associated with *Pycnodonte (Phygraea) vesicularis* (Lamarck) from the Upper Cretaceous of Patagonia. A case of commensalism? *Ameghiniana* 51, 129–140.
- Brezina, S.S., Romero, M.V., Casadío, S., 2017. Encrusting and boring barnacles through the Cretaceous/Paleogene boundary in Northern Patagonia (Argentina). *Ameghiniana* 54, 107–123.
- Busk, G., 1854. Catalogue of the Marine Polyzoa in the Collection of the British Museum. Part 2, Cheilostomata (part). British Museum, London, pp. 55–120.
- Canu, F., 1904. Les bryozoaires du Patagonien. Échelle des bryozoaires pour les terrains tertiaires. *Mémoires de la Société Géologique de France. Paléontologie* 12, 1–30.
- Canu, F., 1911. Iconographie des Bryozoaires fossiles de l'Argentine. Deuxième Partie. *Anales del Museo Nacional de Historia Natural de Buenos Aires* 14, 215–288 (Series 3).
- Canu, F., 1918. Les ovicelles des bryozoaires cyclostomes. Études sur quelques familles nouvelles et anciennes. *Bulletin de la Société Géologique de France* 16, 324–335 (Série 4).
- Canu, F., 1922. Bryozoaires. *Annales de Paléontologie* 11, 124–138.
- Canu, F., Bassler, R.S., 1920. North American Early Tertiary Bryozoa. *Bulletin of the United States National Museum* 106, 1–879.
- Canu, F., Bassler, R.S., 1926. *Phylum Molluscoidea*, vol. 137. United States Geological Survey Professional Paper, pp. 32–39.
- Canu, F., Bassler, R.S., 1929. Bryozoa of the Philippine region. *Bulletin of the United States National Museum* 100 (9), 1–685.
- Carrera, M.G., Casadío, S., 2016. Solenoporacean red algae from lower Danian (Paleogene) coral reefs of northern Patagonia, La Pampa, Argentina. *Ameghiniana* 53, 623–628.
- Casadío, S., 1998. Las ostras del límite Cretácico-Paleógeno de la cuenca Neuquina (Argentina). Su importancia bioestratigráfica y paleobiogeográfica. *Ameghiniana* 35, 449–471.
- Casadío, S., Griffin, M., Parras, A., 2005. *Plicatula* and *Camptonectes* (Bivalvia: Pteriomorpha) from the upper Maastrichtian of Northern Patagonia: Palaeobiogeographic implications. *Cretaceous Research* 26, 507–524.
- Cook, P.L., 1977. The genus *Tremogasterina* Canu (Bryozoa, Cheilostomata). *Bulletin of the British Museum (Natural History), Zoology Series* 35, 103–165.
- Cook, P.L., Bock, P.E., Hayward, P.J., Gordon, D.P., 2018. Class Gymnolaeamata, order Cheilostomata. In: Cook, P.L., Bock, P.E., Gordon, D.P., Weaver, H. (Eds.), *Australian Bryozoa. Volume 2: Taxonomy of Australian families*. CSIRO Publishing, Melbourne, pp. 61–280.
- d'Orbigny, A., 1849–1852. *Prodrome de Paléontologie Stratigraphique Universelle des Animaux Mollusques et Rayonnés faisant suite au cours élémentaire de paléontologie et de géologie stratigraphiques*. Masson, Paris. Vol. 1 (1849), Vol. 2 (1850), Vol. 3 (1852).
- d'Orbigny, A., 1851–1854. *Paléontologie française. Description zoologique et géologique de tous les animaux mollusques et rayonnés fossiles de France, comprenant leur application à la reconnaissance des couches. Terrains crétaçés. Tome cinquième, contenant les bryozoaires*, Victor Masson, Paris, 1192 pp. [1–188

- (1851), 189–472 (1852), 473–984 (1853), 985–1192 (1854)]. Victor Masson, Paris, p. 1192.
- Davis, A.G., 1934. English Lutetian Polyzoa. Proceedings of the Geologists' Association 45, 205–244.
- Di Martino, E., Taylor, P.D., 2013. First bryozoan fauna from a tropical Cretaceous carbonate: Simsim Formation, United Arab Emirates–Oman border region. *Cretaceous Research* 43, 80–96.
- Di Martino, E., Martha, S.O., Taylor, P.D., 2018. The Madagascan Maastrichtian bryozoans of Ferdinand Canu – Systematic revision and scanning electron microscopic study. *Annales de Paléontologie* 104, 101–128.
- Ellis, J., Solander, D., 1786. The natural history of many curious and uncommon zoophytes, collected from various parts of the globe by the late John Ellis. In: Systematically arranged and described by the late Daniel Solander, M.D., F.R.S. Printed for Benjamin White and Son, at Horace's Head, Fleet-Street and Peter Elmsly, in the Strand, London pp. [i]-xii, [1]-208, 63 pls.
- Favorskaya, T.A., 1985. Some species of the genus *Onychocella* from the Maastrichtian of eastern Turkmenistan and western Uzbekistan. *Ezhegodnik Vsesoyuznogo Paleontologicheskogo Obshchestva* 28, 19–33.
- Favorskaya, T.A., 1987. Maastrichtian bryozoans of eastern Turkmeniya. *Ezhegodnik Vsesoyuznogo Paleontologicheskogo Obshchestva* 30, 82–107.
- Favorskaya, T.A., 1988. Maastrichtian bryozoans from western Uzbekistan. Bulletin of the Moscow Society of Naturalists. Department of Geology 63, 94–102.
- Favorskaya, T.A., 1992. Campanian and Maastrichtian bryozoans of the southern USSR. In: Okuneva, T.M., Titova, M.V., Favorskaya, T.A., Zonova, T.D., Rostovtsev, K.O. (Eds.), Atlas rukovodnyashchikh grupp fauny mezozoya Yuga i Vostoka SSSR. Trudy Vserossiyskogo Nauchno-issledovatel'skogo Geologicheskogo Instituta. Nedra, St Petersburg, pp. 115–136.
- Fleming, J., 1828. A history of British animals, exhibiting their descriptive characters and systematical arrangement of the genera and species of quadrupeds, birds, reptiles, fishes, Mollusca, and Radiata of the United Kingdom. Bell and Bradfute, Edinburgh, p. 565.
- Goldfuss, G.A., 1826–1833. *Petrefacta Germaniae*. Teil 1, p. 76. Dusseldorf.
- Gordon, D.P., 1986. The marine fauna of New Zealand: Bryozoa: Gymnolaemata (Ctenostomata and Cheilostomata Anasca) from the western South Island continental shelf and slope, vol. 95. New Zealand Oceanographic Institute Memoir, pp. 1–121.
- Gray, J.E., 1848. List of the Specimens of British Animals in the Collection of the British Museum. Part I. Centroniae or Radiated Animals. Trustees of the British Museum, London xiii + 173.
- Gregory, J.W., 1909. Catalogue of the Fossil Bryozoa in the Department of Geology, British Museum (Natural History). The Cretaceous Bryozoa, vol. 2. Trustees of the British Museum (Natural History), London, p. 346.
- Griffin, M., Casadio, S., Parras, A., 2005. Maastrichtian and Danian species of *Turkostreini* (Ostreidae, Crossostreinae) from the Neuquén Basin, Argentina. *Ameghiniana* 42, 257–276.
- Guha, A.K., Nathan, D.S., 1996. Bryozoan fauna of the Ariyalur Group (Late Cretaceous). Tamilnadu and Pondicherry, India. *Palaeontologia Indica* 49, 1–217 (new series).
- Guler, M.V., González Estebenet, M.S., Navarro, E.L., Astini, R.A., Pérez Panera, J.P., Ottone, E.G., Pileri, D., Paolillo, M.A., 2019. Maastrichtian to Danian Atlantic transgression in the north of Patagonia: A dinoflagellate cyst approach. *Journal of South American Earth Sciences* 92, 552–564.
- Hagenow, F. von, 1839. Monographie der Rügen'schen Kreide-Versteinerungen. Abt. 1. Phytolithen und Polyparien. Neues Jahrbuch für Mineralogie 252–296.
- Hara, U., 2001. Bryozoans from the Eocene of Seymour Island, Antarctic Peninsula. *Palaeontologia Polonica* 60, 33–156.
- Harmelin, J.G., 1976. Le sous-ordre des Tubuliporina (Bryozoaires Cyclostomes) in Méditerranée. *Écologie et systématique*. Mémoires de l'Institut Oceanographique (Monaco) 10, 1–326.
- Harmer, S.F., 1915. Polyzoa of the Siboga Expedition. Part 1. Entoprocta, Ctenostomata and Cyclostomata. Siboga Expedition Reports 28a, 1–180.
- Harmer, S.F., 1926. The Polyzoa of the Siboga Expedition. Part II. Cheilostomata Anasca 28b, 183–501.
- Hincks, T., 1881. Contributions towards a general history of the marine Polyzoa. Part VIII. Foreign Cheilostomata (miscellaneous). *Annals and Magazine of Natural History* 5, 132–136.
- Ihering, H. Von, 1902. Historia de las ostras argentinas. *Anales del Museo Nacional de Historia Natural* 2, 109–125.
- Ihering, H. Von, 1903. Les mollusques des Terrains Crétaciques Supérieurs de l'Argentine Orientale. *Anales del Museo Nacional de Buenos Aires* 9, 193–229 (serie 3, tomo2).
- Ihering, H. Von, 1907. Les Mollusques fossils du Tertiaire et du Crétacé supérieur de l'Argentine. *Anales del Museo Nacional de Buenos Aires* 7, 1–611 (serie 3).
- Illies, G., 1976. Budding and branching patterns in the genera *Stomatopora* Bronn, 1825 and *Voigtapora* Bassler, 1952 (Bryozoa Cyclostomata). *Oberrheinische Geologische Abhandlungen* 25, 97–110.
- Keller, G., Adatte, T., Tantawy, A.A., Berner, Z., Stinnesbeck, W., Stueben, D., Leanza, H.A., 2007. High stress upper Maastrichtian-lower Danian palaeoenvironment in the Neuquén Basin, Argentina. *Cretaceous Research* 28, 939–960.
- Kiessling, W., Aragón, E., Scasso, R., Aberhan, M., Kriwet, J., Medina, F., Fracchia, D., 2005. Massive corals in Paleocene siliciclastic sediments of Chubut (Argentina). *Facies* 51, 233–241.
- Kiessling, W., Scasso, R., Aberhan, M., Ruiz, L., Weidemeyer, S., 2006. A Maastrichtian microbial reef and associated limestones in the Roca Formation of Patagonia (Neuquén Province, Argentina). *Fossil Record* 9 (2), 183–197.
- Koch, C.L., Dunker, W.B.R.H., 1837. Beiträge zur Kenntniss des Norddeutschen Oolithgebildes und dessen Versteinerungen. Tome 4, 1–64.
- Koromysova, A.V., Seltser, V.B., 2020. Early Maastrichtian cheilostome bryozoans from the middle Volga River region. *PalZ* 94, 697–714.
- Koromysova, A.V., Baraboshkin, E.Y., Martha, S.O., 2018a. Late Campanian to upper Maastrichtian bryozoans encrusting on belemnite rostra from the Aktolagay Plateau in western Kazakhstan. *Geobios* 51, 307–333.
- Koromysova, A.V., Martha, S.O., Pakhnevich, A.V., 2018b. The internal morphology of *Acoscinopectora* Voigt, 1956 (Cheilostomata, Bryozoa) from the Campanian–Maastrichtian of Central and Eastern Europe. *PalZ* 92, 241–266.
- Koromysova, A.V., Taylor, P.D., Martha, S.O., Riley, M., 2018c. *Rhagasostoma* (Bryozoa) from the Late Cretaceous of Eurasia: taxonomic revision, stratigraphy and palaeobiogeography. *European Journal of Taxonomy* 490, 1–66.
- Koromysova, A.V., Martha, S.O., Pakhnevich, A.V., 2019a. Revision of *Porina*-like cheilostome Bryozoa from the Campanian and Maastrichtian of Central Asia. *Annales de Paléontologie* 105, 1–19.
- Koromysova, A.V., Pakhnevich, A.V., Fedorov, P.V., 2019b. *Tobolocella levinae* n. gen., n. sp., a cheilostome bryozoan from the upper Maastrichtian of Northern Kazakhstan: Scanning electron microscope and micro-CT study. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 294, 91–101.
- Lamarck, J.B.P.A. de, 1806. Sur les fossiles des environs de Paris. Genre 74 Huître Ostrea. *Annales Muséum Histoire Naturelle* 8, 156–166.
- Lamarck, J.B.P.A. de, 1816. Les Polypes. *Histoire naturelle des Animaux sans Vertèbre* II. Paris, p. 586.
- Lamouroux, J.V.F., 1816. *Histoire des Polypiers Coralligènes Flexibles*, vulgairement nommés Zoophytes. Caen, lxxiv + 559.
- Lang, W.D., 1916. A revision of the "cribrimorph" Cretaceous Polyzoa. *Annals and Magazine of Natural History (Series 8)* 18 (81–112), 381–410.
- Levinson, G.M.R., 1909. Morphological and systematic studies on the cheilostomatous Bryozoa. *Nationale Forfatterers Forlag, Copenhagen*, pp. 1–431.
- Linnaeus, C., 1767. *Systema Naturae* (12th Edition), vol. 1. *Regnum Animale*. Part 2. *Laurentii Salvii, Holmiae*, pp. 533–1327.
- Malumián, N., Nález, C., 2011. The Late Cretaceous–Cenozoic transgressions in Patagonia and the Fuegian Andes: foraminifera, palaeoecology, and palaeogeography. *Biological Journal of the Linnean Society* 103, 269–288.
- Marsson, T., 1887. Die Bryozoen der weissen Schreiekreide der Insel Rügen. *Palaeontologische Abhandlungen* 4, 1–112.
- Nikulina, E.A., 2007. *Einhornia*, a new genus for electrids formerly classified as the *Electra crustulenta* species group (Bryozoa, Cheilostomata). *Schriften des Naturwissenschaftlichen Vereins für Schleswig-Holstein* 69, 29–40.
- Nikulina, E.A., de Blauwe, H., Reverter-Gil, O., 2012. Molecular phylogenetic analysis confirms the species status of *Electra verticillata* (Ellis and Solander, 1786). In: Ernst, A., Schäfer, P., Scholz, J. (Eds.), *Bryozoa Studies 2010*. Springer, Heidelberg, pp. 217–236.
- Nye jr., O.B., 1976. Generic revision and skeletal morphology of some cerioporid cyclostomes (Bryozoa). *Bulletins of American Paleontology* 69, 1–222.
- Pitt, L.J., Taylor, P.D., 1990. Cretaceous Bryozoa from the Faringdon Sponge Gravel (Aptian) of Oxfordshire. *Bulletin of the British Museum (Natural History), Geology Series* 46, 61–152.
- Prámparo, M.B., Cione, A.L., Gonzalez Riga, B., 2014. Sharks (Neoselachii) and palynomorphs from Mendoza (Argentina): New evidence of the Late Cretaceous Atlantic marine transgression. *Alcheringa* 38, 177–189.
- Römer, F.A., 1840. Die Versteinerungen des norddeutschen Kreidegebirges. *Hahn'schen Hofbuchhandlung, Hannover*, p. 145.
- Scasso, R.A., Concheyro, A., Kiessling, W., Aberhan, M., Hecht, L., Medina, F.A., Tagle, R., 2005. A tsunami deposit at the Cretaceous/Paleogene boundary in the Neuquén Basin of Argentina. *Cretaceous Research* 26, 283–297.
- Shaw, N.G., 1967. Cheilostomata from Gulf (Upper Cretaceous) rocks of southwestern Arkansas. *Journal of Paleontology* 41, 1393–1432.
- Sogot, C.E., Harper, E.M., Taylor, P.D., 2013. Biogeographical and ecological patterns in bryozoans across the Cretaceous–Paleogene boundary: Implications for the phytoplankton collapse hypothesis. *Geology* 41, 631–634.
- Sogot, C.E., Harper, E.M., Taylor, P.D., 2014. The Lilliput effect in colonial organisms: Cheilostome bryozoans at the Cretaceous–Paleogene mass extinction. *PLoS One* 9, e87048.
- Stilwell, J.D., Håkansson, E., 2012. Survival, but...! New tales of 'Dead Clade Walking' from Austral and Boreal post-K-T assemblages. In: Talent, J.A. (Ed.), *Earth and Life. Global Biodiversity, Extinction Intervals and Biogeographic Perturbations Through Time*. Springer, Dordrecht, pp. 795–810.
- Taylor, P.D., 2008. Late Cretaceous cheilostome bryozoans from California and Baja California. *Journal of Paleontology* 82, 823–834.
- Taylor, P.D., 2019. A brief review of the scanty fossil record of Cretaceous bryozoans from Gondwana. *Australasian Palaeontological Memoirs* 52, 147–154.
- Taylor, P.D., Brezina, S.S., 2018. A new Cenozoic cyclostome bryozoan genus from Argentina and New Zealand: Strengthening the biogeographical links between South America and Australasia. *Alcheringa* 42, 441–446.
- Taylor, P.D., Di Martino, E., 2018. *Sonarina tamilensis* n. gen., n. sp., an unusual cheilostome bryozoan from the Late Cretaceous of southern India. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 288, 79–85.

- Taylor, P.D., McKinney, F.K., 2006. Cretaceous Bryozoa from the Campanian and Maastrichtian of the Atlantic Coastal Plains, United States. *Scripta Geologica* 132, 1–346.
- Taylor, P.D., Sequeiros, L., 1982. Toarcian bryozoans from Belchite in north-east Spain. *Bulletin of the British Museum (Natural History), Geology Series* 36, 117–129.
- Taylor, P.D., Waeschenbach, A., 2015. Phylogeny and diversification of bryozoans. *Palaeontology* 58, 585–599.
- Taylor, P.D., Wilson, M.A., 2003. Palaeoecology and evolution of marine hard substrate communities. *Earth-Science Reviews* 62, 1–103.
- Taylor, P.D., Zatoń, M., 2008. Taxonomy of the bryozoan genera *Oncousoecia*, *Microeciella* and *Eurystrotos* (Cyclostomata: Oncousoeciidae). *Journal of Natural History* 42, 2557–2574.
- Taylor, P.D., Di Martino, E., Martha, S.O., 2018. Colony growth strategies, dormancy and repair in some Late Cretaceous encrusting bryozoans: insights into the ecology of the Chalk seabed. *Palaeobiodiversity and Palaeoenvironments* 99, 425–446.
- Turner, R.F., 1979. North American Cretaceous Tricéphaloporinae. In: Larwood, G.P., Abbott, M.B. (Eds.), *Advances in Bryozoology*. Academic Press, London, pp. 521–539.
- Uttley, G.H., 1951. The Recent and Tertiary Polyzoa (Bryozoa) in the collection of the Canterbury Museum, Christchurch. Part II. Records of the Canterbury Museum 6, 15–39.
- Voigt, E., 1967. Oberkreide-Bryozoen aus den asiatischen Gebieten der UdSSR. *Mitteilungen aus dem Geologischen Staatsinstitut in Hamburg* 36, 5–95.
- Walter, B., 1970. Les Bryozoaires Jurassiques en France, vol. 35. Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon, pp. 1–328 [for 1969].
- de Winter, N.J., Vellekoop, J., Vorsselmans, R., Golreihan, A., Soete, J., Petersen, S.V., Meyer, W., Casadio, S., Speijer, R.P., Claeys, P., 2018. An assessment of latest Cretaceous *Pycnodonte vesicularis* (Lamarck, 1806) shells as records for palaeoseasonality: a multi-proxy investigation. *Climate of the Past* 14, 725–749.
- Woelders, L., Vellekoop, J., Kroon, D., Smit, J., Casadio, S., Prámparo, M.B., Dinarès-Turell, J., Peterse, F., Sluijs, A., Lenaerts, J.T.M., Speijer, R.P., 2017. Latest Cretaceous climatic and environmental change in the South Atlantic region. *Paleoceanography* 32, 466–483.