

**NEW RECORDS OF VERTEBRATES AND TRACE FOSSILS FROM THE HUINCUL FORMATION  
(CENOMANIAN-TURONIAN, UPPER CRETACEOUS) IN THE VICINITY OF EL CHOCÓN, NEUQUÉN  
PROVINCE, ARGENTINA**

J. I. CANALE<sup>1,2,3</sup>, A. C. GARRIDO<sup>4</sup>, P. CITTON<sup>1,5</sup>, S. DE VALAIS<sup>1,5</sup>, M. A. BAIANO<sup>1,2,3</sup>, D. PALOMBI<sup>1,2</sup>, M. CARATELLI<sup>1,5</sup>, S. APESTEGUÍA<sup>1,6</sup>, I. MANIEL<sup>1,7</sup> AND M. L. FERNANDEZ DUMONT<sup>1,6\*</sup>

<sup>1</sup>Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

<sup>2</sup>Área Laboratorio e Investigación, Museo Municipal "Ernesto Bachmann". Dr. Natali s/n, Q8311AZA Villa El Chocón, Neuquén, Argentina.  
*jicanale@unrn.edu.ar; mbaiano@unrn.edu.ar*

<sup>3</sup>Universidad Nacional de Río Negro (UNRN). Isidro Lobo 516, 8332 General Roca, Río Negro, Argentina.

<sup>4</sup>Museo Provincial de Ciencias Naturales "Prof. Dr. Juan Olsacher". Zapala, Neuquén, Argentina.

<sup>5</sup>Instituto de Investigación en Paleobiología y Geología (IIPG, UNRN). Av. Roca 1242, 8332 General Roca, Río Negro, Argentina.  
*pcitton@unrn.edu.ar; sdevalais@yahoo.com.ar; mcaratelli@unrn.edu.ar*

<sup>6</sup>Fundación de Historia Natural Félix de Azara. Centro de Ciencias Naturales Ambientales y Antropológicas, Universidad Maimónides. Hidalgo 775, 7mo piso, 1405 Ciudad Autónoma de Buenos Aires, Argentina. *mlucilafd@gmail.com*

<sup>7</sup>Instituto de Evolución, Ecología Histórica y Ambiente (UTN). Urquiza 314, 5600 San Rafael, Mendoza, Argentina.  
*nachomaniel@gmail.com*

The Huincul Formation (Neuquén Group) has preserved a rich fauna of continental fossil vertebrates, biased towards large forms, almost exclusively dinosaurs. This contrasts with the underlying Candeleros Formation, where records of small vertebrates and tracks abound. Here we present new trace and corporeous fossils records from the lower section of the Huincul Formation, outcropping at the Cañadón del Cocodrilo site, about 4 km NE of Villa El Chocón. Sub-circular to ovoidal depressions of decimetric dimension are preserved on the top-surface of reddish, fine- to medium-grained sandstones. Concave hyporeliefes show mutual alignment and narrow, fairly developed rims bordering their outer edge, suggesting identifying them as dinosaur tracks. A single kidney-shaped hyporelief, resembling a sauropod manus print in morphology, further supports our hypothesis. Invertebrate traces are straight to gently sinuous and variably oriented, unlined, cylindrical, meniscate backfilled burrows, up to 18 mm wide, with rare secondary branching. Invertebrate traces resemble the ichnospecies *Taenidium barretti*, indicative of the 'pre-desiccation' suite of the *Scyenia ichnofacies*. The vertebrate remains collected include part of a hypoplastron (Museo Municipal "Ernesto Bachmann" collection: MMCh-PV 302), which on the basis of the ornamentation and the loose connection of the inguinal pillar between the carapace and the plastron, is referred to a turtle Chelidae pleurodiran. It is remarkable that the position of the abdominal-femoral sulcus is different to that present in other fossil chelids of the Neuquina Basin. A fragment of a right mandible (MMCh-PV 303) was collected, whose densely packed acrodont teeth, with a subtriangular section, allow its assignment to a sphenodontian lepidosaur Eilenodontinae, cf. *Priosphenodon* sp. A partial skeleton (MMCh-PV 304) was recovered including an incomplete skull which displays subcircular supratemporal fenestrae and unornamented quadratojugal, that, along with the small size of the specimen allow to assign it to the crocodyliform Notosuchia Uruguaysuchidae. Some features, such as the narrow dorsal surface of the parietal between the supratemporal fenestrae, elevated rims along the margins of these fenestrae, the low medial longitudinal crest over the frontals and angular with and oblique and strongly marked crest, resemble the species *Araripesuchus buitreraensis*. This is the first record of vertebrate tracks, chelid turtle, and notosuchian crocodile remains at the Huincul Formation. This assemblage strongly expands the knowledge of the fossil fauna for this geological unit, and helps to fill the record gap of certain groups of vertebrates in the Patagonian Cretaceous.

\*Financial support provided by: PICT 2018-04042 and Municipalidad de Villa El Chocón.